

On AGT relation in the case of $U(3)$

A.Mironov* and A.Morozov†

Abstract

We consider the AGT relation [1], expressing conformal blocks for the Virasoro and W -algebras in terms of Nekrasov's special functions, in the simplest case of the 4-point functions for the first non-trivial W_3 algebra. The standard set of Nekrasov functions is sufficient only if additional null-vector restriction is imposed on a half of the external W -primaries and this is just the case when the conformal blocks are fully dictated by W -symmetry and do not depend on a particular model. Explicit checks confirm that the AGT relation survives in this restricted case, as expected.

Contents

1	Introduction	2
2	The simplest example: two free fields with $c = 2$	3
2.1	Structure of the W_3 algebra	3
2.2	4-point conformal block for the W_3 algebra	4
2.3	Nekrasov's formulas	5
2.4	AGT relation	5
2.4.1	Denominator	6
2.4.2	Numerator. The case of $\vec{\alpha}_1 = \dots = \vec{\alpha}_4 = 0$	6
2.4.3	Numerator. The case of arbitrary $\vec{\alpha}_2$ and $\vec{\alpha}_4$ with no ambiguity in the W -conformal block	7
2.4.4	The case of $\xi = 2$	8
2.4.5	The $U(1)$ factor ν and projective transformations	8
2.4.6	General case, all $\vec{\alpha} \neq 0$, $\vec{\alpha}_1$ and $\vec{\alpha}_3$ <i>special</i>	8
2.5	Intermediate summary	10
3	Some CFT considerations	10
3.1	W -algebra and Shapovalov form	10
3.2	3-point functions at level one	11
3.3	Shapovalov matrix at level two	12
4	AGT relation to $c \neq 2$ at level one	14
4.1	No external lines, $\vec{\alpha}_1, \dots, \vec{\alpha}_4 = 0$	14
4.2	Introducing external states	15
5	Calculations at level two	16
5.1	Level-two Nekrasov functions	16
5.2	AGT relations at level two	17
5.3	Factorization of Kac determinant	18
5.4	Conformal block at level two	19
6	On complete proof of AGT relation in a <i>very special</i> case	21
7	Summary and conclusion	23

*Lebedev Physics Institute and ITEP, Moscow, Russia; mironov@itep.ru; mironov@lpi.ru

†ITEP, Moscow, Russia; morozov@itep.ru

1 Introduction

A recent paper [1], see also [2]-[6], provided a long-anticipated relation between $2d$ conformal theories [7] and Nekrasov's functions [8], decomposing the Seiberg-Witten prepotential into sums over $4d$ instantons. An exact formulation of the AGT relation [1] is that the **universal conformal blocks [7, 9, 10] for generic Verma modules of the Virasoro algebra coincide with the Nekrasov partition functions, provided the α -parameters of the Virasoro primary fields are linear combinations of the a and μ -parameters of the Nekrasov functions.** The central charge is related to Nekrasov's peculiar additional parameter $\epsilon = \epsilon_1 + \epsilon_2$, $c = 1 + \frac{6\epsilon^2}{\epsilon_1\epsilon_2}$ and dimensions are given by $\Delta_\alpha = \frac{\alpha(\epsilon-\alpha)}{\epsilon_1\epsilon_2}$. For details of explicit check of this relation see [1, 2, 4]. An additional observation of [1] is that with the same $\{\alpha\} - \{a, \mu\}$ identification of the perturbative part of the Nekrasov partition function (exponentiated perturbative part of the Seiberg-Witten prepotential) coincides with the DOZZ functions [11]: the structure constants of concrete, Liouville conformal model, that is, a certain reduction of the $SU(2)$ WZNW model, which has a natural description in terms of a single $2d$ free field [12]. With no surprise Nekrasov's functions, appearing in this original AGT relation are associated with the gauge group $SU(2)$.

An obvious generalization [2] of the AGT relation is to take a $2d$ conformal model made from the $r = N - 1$ free fields with the extended W_N -symmetry [13]-[17] and look if its conformal blocks are expanded into the $SU(N)$ Nekrasov functions.

The first problem here is that the theory of W -symmetries is under-developed. In the Virasoro sector, the crucial idea is to identify Virasoro operators with the $2d$ stress-tensor, which generates shifts in the holomorphic z coordinate. This identification results in the two well-known postulates:

$$L_{-1}V(z) = \partial_z V(z) \quad (1)$$

for arbitrary vertex operator $V(z)$ and

$$\langle V_{\Delta_1}(z_1)V_{\Delta_2}(z_2)V_{\Delta_3}(z_3) \rangle \sim z_{12}^{\Delta_3-\Delta_1-\Delta_2} z_{23}^{\Delta_1-\Delta_2-\Delta_3} z_{31}^{\Delta_2-\Delta_1-\Delta_3} \quad (2)$$

for an arbitrary spherical 3-point function of the Virasoro *quasi primaries* [7]. The 3-point correlator of secondary fields can be then immediately constructed from this one by exploiting the Virasoro symmetry. After that, arbitrary conformal blocks and correlators can be built from these three point functions by a "gluing-of-pants" procedure, at least in principle [7, 18, 19]. Though it is a difficult procedure, involving infinite summations, it is unambiguously defined. In the case of W -symmetries, there is still no clear choice for the W -generalization of z (a modulus of the W -geometry, see, e.g., [20]) and, hence, no formulas like (1) and (2). In result, an arbitrary theory with W -symmetry has many additional free parameters: the entire set of the 3-point functions in the W_3 case is

$$\langle V_{\vec{\alpha}_1}(1) V_{\vec{\alpha}_2}(\infty) (W_{-1})^k V_{\vec{\alpha}}(0) \rangle \quad (3)$$

involving a single insertion of arbitrary power of W_{-1} . In a theory with underlying W -geometry, like the conformal Toda proposed as an obvious $SU(N)$ -counterpart of the Liouville theory in [2], these infinitely many parameters are of course fixed, but the relevant theory is still unavailable.

Another obvious problem concerns the AGT relations themselves. For $N > 2$ there is a clear mismatch in the number of parameters. Each W_N primary, external or internal, is labeled by the highest-weight, which is arbitrary $r = N - 1$ dimensional vector $\vec{\alpha}$, thus the 4-point conformal block depends on $(4+1) \cdot (N-1) + 1 = 5N - 4$ free parameters (4 + 1 is the number of external + internal states, an extra 1 is the central charge). At the same time, Nekrasov function depends on $N - 1$ "expectation values" $\vec{a}$, $2N$ "masses" μ_f and two ϵ -parameters, what gives $(N - 1) + 2N + 2 - 1 = 3N$ (1 is subtracted because conformal invariance forces Nekrasov functions to be dimensionless: to depend only on ratios of parameters). For $N = 2$ there was a nice matching [4]: $5N - 4 = 3N$, but it breaks down for $N > 2$: some $2(N - 2)$ parameters of the conformal block do not have immediate counterparts in the Nekrasov functions.

These two difficulties could be considered as a strong appeal for developing the theory of W -symmetries and looking for a further generalization of Nekrasov functions. However, Niclas Wyllard suggested in [2] an elegant way out which overcomes both problems at once. One can restrict consideration to the case when the appropriate number of "external" states are the *special* ones [17], satisfying additional W -null-vector constraints at level one¹. In the case of W_3 just one is enough,

$$(3wL_{-1} - 2\Delta W_{-1})V_{\Delta,w} = 0, \quad (4)$$

¹In fact, restriction to the special states is rather natural within Davide Gaiotto's construction [21] behind the AGT relation, where $4d$ SYM theory is described as a result of 5-brane compactification on a Riemann surface (spectral curve) *a la* [22, 23]. For $N \geq 3$ the punctures on the Riemann surface are not of general type, and it was suggested in [2] to identify them exactly with the *special* states on the CFT side of the AGT relation. We are grateful to the referee of the paper for his comments on this point.

for higher N the number of constraints grows up to $N-2$.² Such "special" primary states form a one-parametric family instead of an r -parametric one. "The appropriate number" to solve the matching problem is two out of four external states for the 4-point conformal block. At the same time, if one of the three primaries in (3) is special, these 3-point functions get unambiguously defined without any reference to W -geometry. Thus, taking two of external states to be *special*, one also eliminates the first problem. Moreover, just under this restriction a counterpart of the DOZZ functions for the $SU(3)$ conformal Toda has been evaluated in [17], and [2] checks that they indeed coincide with the exponential of perturbative $SU(3)$ prepotential in the conformal invariant case $N_f = 2N = 6$. In particular, the linear relation between α and a, μ parameters was established in this case.

Unfortunately no detailed check of the main part of the AGT relation, between conformal blocks and "instanton sums", was explicitly made in [2], despite relevant formulas were extracted from [13]-[17]. It is our task in this paper to partly fill in this gap. We explicitly present this check in the simplest case. The "simplest" involves the following restrictions: only $N = 3$ and only contributions at levels one and two are considered. It is not a big problem to lift any of these two restrictions, still calculations are tedious, and it is unclear if they are that necessary to provide even more evidence in support of the AGT relation. In any case, they remain for the future work, maybe they can help to better understand the notion of W -symmetry beyond the seemingly-artificial (4) constraint and to reveal the true meaning of the AGT relations, and the way to prove it in the full generality. This would further promote Nekrasov functions in the role of the crucially important new special functions and stimulate the further study of matrix-model tau-functions (i.e. generalized τ -functions [24], subjected to additional Virasoro-like constraints), already proposed for this role in [25].

Presentation in this paper does *not* follow literally the by-now-standard one, accepted in [1, 2, 4]. Instead of just writing down the conformal blocks and Nekrasov functions and comparing them (actually reporting the result of computer calculations), we *also* try to demonstrate that the AGT relation itself can turn into a powerful alternative method of iterative, level-by-level, evaluation of conformal blocks. Therefore, we begin with making ansatzes for the elementary constituents of conformal blocks (the 2- and 3-point functions) and then define the remaining coefficients by the requirement that the result matches Nekrasov functions. A part of this comparison is to check that the Kac determinant (that of the Shapovalov matrix made from the scalar products of states, essentially a 2-point functions at special points $z_1 = 0, z_2 = \infty$) factorizes in an appropriate way: in α -parametrization of primary fields its roots are given by linear function $\vec{\alpha}\vec{e}_i = m_i\epsilon_1 + n_i\epsilon_2$. The AGT relation strengthens this well known Regge-trajectory-like statement and makes an additional claim about the numerators of conformal blocks, not only denominators: **the Nekrasov expansion provides a parametrization of the numerators by μ -variables, which are also linear functions of α -parameters.**

Only after making this kind of presentation at level one and for the simplest case of $c = 2$ in s.2, we turn to the standard approach: calculate conformal blocks by the standard CFT methods in s.3, what also justifies the results of s.2. The necessary Ward identities, providing recursive relations for the 3-point functions, are taken from a separate summary in [5]. Then in s.4 we extend the check of the $U(3)$ AGT relation to $c \neq 2$ and in s.5 to level two. At last, in s.6 we provide the **complete** proof of the AGT relation in the very particular case of one of the special states in the conformal block being completely degenerate at the first level. The proof is possible, since there is a complete answer for the conformal block in this case [17].

2 The simplest example: two free fields with $c = 2$

2.1 Structure of the W_3 algebra

In the case of several free fields the set of Virasoro primaries is not exhausted by exponentials $e^{\sqrt{2}\vec{\alpha}\vec{\phi}}$: there are many more primaries. If one wants exponentials to remain the only primary fields (in addition to the currents $\partial\vec{\phi}$), they should be primaries of a larger algebra, extending the Virasoro one. The standard choice in the case of r free fields is the W_{r+1} algebra.

Generators $W^{(k)}(z) = \omega^{i_1 \dots i_k} \partial\phi_{i_1} \dots \partial\phi_{i_k}(z)$ and tensors ω are defined so that only minimal singularities are allowed in the algebra. From now on, we switch to the simplest example of two free fields, $r = 2$, associated with the $SU(3)$ group and W_3 algebra. The W_3 algebra has two generators: the stress tensor $T(z) = W^{(2)}(z) =$

In the present text, we discuss a possibility to express conformal blocks through Nekrasov functions *without* explicit reference to branes or to instanton expansions in Seiberg-Witten theory, i.e. we are using AGT relations as a tool to find explicit expressions for conformal blocks, and then restriction to the *special* states should be eliminated one day, perhaps, by further extending the set of Nekrasov functions.

²To avoid possible confusion, note that in [17, (2.12)] some three constraints were imposed on one of the special states. This was, however, done to make a *particular method* working, as to the conformal block, it is well defined (but very hard to evaluate) already with a single constraint for the special state.

$\frac{1}{2}(\partial\vec{\phi})^2(z)$ and $W(z) = W^{(3)}(z)$. The stress tensor $T(z)$ is invariant under $SO(2)$ rotations of ϕ , and this rotation freedom should be fixed in order to define $W(z)$ unambiguously. We require that it is symmetric under $\phi_2 \rightarrow -\phi_2$ and, therefore, antisymmetric under $\phi_1 \rightarrow -\phi_1$. This means that $W = (\partial\phi_1)^3 + h\partial\phi_1(\partial\phi_2)^2$ with the single undefined parameter h . The operator product expansion is

$$T(z)W(0) = \frac{3+h}{z_4}\partial\phi_1(0) + \dots \quad (5)$$

and h is fixed by the requirement of absence the most singular term with z^{-4} (which is equivalent to the requiring the W -field to be Virasoro primary). This defines h to be $h = -3$ and

$$W(z) = W^{(3)}(z) \sim (\partial\phi_1)^3 - 3\partial\phi_1(\partial\phi_2)^2 = \partial\phi_1 \left((\partial\phi_1)^2 - 3(\partial\phi_2)^2 \right) \quad (6)$$

(the normalization coefficient is actually $2^{-3/2}$, see the Appendix). Accordingly,

$$\begin{aligned} L_0 e^{\sqrt{2}(\alpha\phi_1 + \beta\phi_2)} &= \Delta_{\alpha,\beta} e^{\sqrt{2}(\alpha\phi_1 + \beta\phi_2)}, \\ W_0 e^{\sqrt{2}(\alpha\phi_1 + \beta\phi_2)} &= w_{\alpha,\beta} e^{\sqrt{2}(\alpha\phi_1 + \beta\phi_2)} \end{aligned} \quad (7)$$

with

$$\begin{aligned} \Delta_{\alpha,\beta} &= \alpha^2 + \beta^2, \\ w_{\alpha,\beta} &= \alpha(\alpha^2 - 3\beta^2) \end{aligned} \quad (8)$$

Crucial for the AGT relation in the $SU(3)$ case is the fact, that

$$\Delta_{\alpha,\beta}^3 - w_{\alpha,\beta}^2 = \left(\alpha^2 + \beta^2 \right)^3 - \left(\alpha(\alpha^2 - 3\beta^2) \right)^2 = \left(\beta(\beta^2 - 3\alpha^2) \right)^2 = w_{\beta,\alpha}^2 \quad (9)$$

The r.h.s. has nothing to do with the W_3 algebra, but this relation demonstrates that a certain linear combination of Δ and w , that is, the one at the l.h.s. is a full square.

2.2 4-point conformal block for the W_3 algebra

As reviewed in [4] and (in far more detail) in [5], the conformal block has the structure

$$B_{\vec{\alpha}_1, \vec{\alpha}_2, \vec{\alpha}_3, \vec{\alpha}_4}^{\vec{\alpha}}(x) = \sum_{|\mathcal{Y}|=|\mathcal{Y}'|} x^{|\mathcal{Y}|} \mathcal{B}_{\vec{\alpha}_1, \vec{\alpha}_2, \vec{\alpha}_3, \vec{\alpha}_4}^{\vec{\alpha}}(\mathcal{Y}, \mathcal{Y}') = \sum_{|\mathcal{Y}|=|\mathcal{Y}'|} x^{|\mathcal{Y}|} \bar{\Gamma}_{\vec{\alpha}_1, \vec{\alpha}_2; \vec{\alpha}}(\mathcal{Y}) Q_{\vec{\alpha}}^{-1}(\mathcal{Y}, \mathcal{Y}') \Gamma_{\vec{\alpha}, \vec{\alpha}_3, \vec{\alpha}_4}(\mathcal{Y}') \quad (10)$$

Here $\mathcal{Y}$ labels the elements of the Verma module for the W_3 algebra, i.e. a generalization to W_3 of what was the Young diagram for the Virasoro algebra, $\bar{\Gamma}(\mathcal{Y})$ and $\Gamma(\mathcal{Y})$ are the two types of the corresponding 3-point functions (they are essentially the same in the Virasoro case, but interrelations are more sophisticated in the W -sector [5]) and $Q(\mathcal{Y}, \mathcal{Y}')$ is the Shapovalov form: a table made from the scalar products of different elements in the Verma module, which, as usual, does not mix different levels $|\mathcal{Y}|$.

At level one, $|\mathcal{Y}| = 1$ there are exactly two states in the Verma module, $L_{-1}V_{\vec{\alpha}}$ and $W_{-1}V_{\vec{\alpha}}$ and the corresponding block of the Shapovalov form is

$Q_{\vec{\alpha}}(\mathcal{Y}, \mathcal{Y}')$	$L_{-1}V_{\vec{\alpha}}$	$W_{-1}V_{\vec{\alpha}}$
$L_{-1}V_{\vec{\alpha}}$	$2\Delta_{\vec{\alpha}}$	$3w_{\vec{\alpha}}$
$W_{-1}V_{\vec{\alpha}}$	$3w_{\vec{\alpha}}$	$q\Delta_{\vec{\alpha}}^2$

(11)

It is straightforwardly evaluated from the commutation relations of the algebra. However, at level one of the W_3 -algebra the entries of this simple matrix are defined on dimensional grounds up to the coefficients. Coefficients 2 and 3 are dictated by the elementary part of W_3 commutators,

$$\begin{aligned} [L_n, L_m] &= (n-m)L_{m+n} + \frac{c}{12}n(n^2-1), \\ [L_n, W_m] &= (2n-m)W_{m+n} \end{aligned} \quad (12)$$

while q follows from the sophisticated one for $[W_m, W_n]$, which also depends on the normalization of $W(z)$. This remaining commutation relation will appear only in (50) far below. Instead of using it now, one can just *guess* that $q = 9/2$ by looking at (9): then the determinant $2q\Delta^3 - 9w^2$ of the Shapovalov form (Kac determinant), which stands in the denominator of conformal block at level one, factorizes nicely: moreover, it is a full square.

For this value of $q = 9/2$

$$B_{\vec{\alpha}_1, \vec{\alpha}_2, \vec{\alpha}_3, \vec{\alpha}_4}^{\vec{\alpha}}(x) = 1 + \frac{xK_1}{\Delta_{\vec{\alpha}}^3 - w_{\vec{\alpha}}^2} + O(x^2) \quad (13)$$

or

$$B_1 = \frac{K_1}{\Delta_{\vec{\alpha}}^3 - w_{\vec{\alpha}}^2} \stackrel{(9)}{=} \frac{K_1}{v^2} \quad (14)$$

where v^2 is the r.h.s. of (9) and all dependencies on $\vec{\alpha}_1, \dots, \alpha_4$ are contained in the numerator K_1 .

2.3 Nekrasov's formulas

Nekrasov's partition function has the form

$$\mathcal{Z} = \sum_{Y_1, \dots, Y_N} x^{|Y_1| + \dots + |Y_N|} \mathcal{Z}(Y_1, \dots, Y_N) \quad (15)$$

where the sum is over N sets of the *ordinary* Young diagrams. The level one contribution to $\mathcal{Z}$ is simply

$$\mathcal{Z} = 1 + x \sum_{i=1}^N \mathcal{Z}(\emptyset, \dots, \square, \dots, \emptyset) + O(x^2) \quad (16)$$

where the only non-empty diagram $Y = \square$ stands at the i -th place. Explicitly, for the conformal invariant case $N_f = 2N$ one has

$$Z_1 = \sum_{i=1}^N \mathcal{Z}(\emptyset, \dots, \square, \dots, \emptyset) = -\frac{1}{\epsilon_1 \epsilon_2} \sum_{i=1}^N \frac{P(a_i)}{\prod_{j \neq i}^N (a_i - a_j)(a_i - a_j + \epsilon)} \quad (17)$$

where $P(a_i) = \prod_{f=1}^{2N} (a_i + \mu_f)$. In our simple example $\epsilon = 0$ and Z_1 has square of the Van-der-Monde determinant $\Delta(\vec{a}) = \prod_{i < j} (a_i - a_j)$ in the denominator,

$$Z_1(\epsilon = 0) = \frac{M_1}{\Delta(\vec{a})^2} \quad (18)$$

We preserve the standard notation Δ for both dimensions and the Van-der-Monde determinants, hopefully this will not cause a confusion. In the dimension, the argument $\vec{\alpha}$ is a subscript, while in the determinant $\vec{a}$ is an argument in brackets. The vector $\vec{a}$ is actually $r = N - 1$ -dimensional, like $\vec{\alpha}$, since its N components a_i (eigenvalues of the gauge-field vev matrix) are constrained by the zero trace condition

$$\sum_{i=1}^N a_i = 0 \quad (19)$$

2.4 AGT relation

The AGT relation [1, 2, 4] is the statement that for some linear relation between the whole set of $\vec{\alpha}$'s and the whole set of $\vec{a}$ and μ_f , the conformal blocks $\mathcal{B}$ for the W_N algebra and the Nekrasov partition function $\mathcal{Z}$ for $U(N)$ group coincide provided $c = (N - 1) \left(1 + N(N + 1) \frac{\epsilon^2}{\epsilon_1 \epsilon_2} \right)$. We start from the simplest version of this relation for $N = 3$: for the 4-point conformal block and for $c = 2$, i.e. $\epsilon = 0$.

Comparing (14) and (18) one observes that in this case the AGT relation requires that

$$v \sim \Delta(\vec{a}) \quad (20)$$

which dictates the relation between $\vec{a}$ and $\vec{\alpha}$ of the intermediate state, and

$$K_1 \sim M_1 \quad (21)$$

which would define the relation between μ_f and $\vec{\alpha}_1, \dots, \vec{\alpha}_4$ for the four external states.

2.4.1 Denominator

Let us begin with (20). Coming back to the parametrization $\vec{\alpha} = (\alpha, \beta)$, one obtains

$$v \stackrel{(9)}{=} \beta(\beta^2 - 3\alpha^2) = 4\beta \cdot \frac{\beta + \alpha\sqrt{3}}{2} \cdot \frac{\beta - \alpha\sqrt{3}}{2} \sim (a_1 - a_2)(a_2 - a_3)(a_3 - a_1) \quad (22)$$

which, together with (19), implies a linear relation of the form

$$\begin{aligned} a_1 &= \kappa(\alpha - \beta\sqrt{3}), \\ a_2 &= \kappa(\alpha + \beta\sqrt{3}), \\ a_3 &= -2\kappa\alpha \end{aligned} \quad (23)$$

It can be, of course, written in terms of the root and weight vectors of $SU(3)$.

2.4.2 Numerator. The case of $\vec{\alpha}_1 = \dots = \vec{\alpha}_4 = 0$

Let us now proceed to (21). We begin with putting all the four "external" dimensions equal to zero, $\alpha_1 = \beta_1 = \dots = \alpha_4 = \beta_4 = 0$. Then (21) follows from

$$B^{(1)} = \frac{\Delta(p\Delta^3 + sw^2)}{\Delta^3 - w^2} = -\frac{1}{\epsilon_1\epsilon_2} \left(\frac{P_6(a_1)}{a_{12}^2 a_{23}^2} + \frac{P_6(a_2)}{a_{21}^2 a_{23}^2} + \frac{P_6(a_3)}{a_{31}^2 a_{32}^2} \right) = Z^{(1)} \quad (24)$$

where only the two parameters p and s at the l.h.s. are not defined on dimensional grounds and $P_6(a) = a^6 + \sum_{k=1}^6 \sigma_k a^{6-k}$ at the r.h.s. is a polynomial of degree 6, depending on choice of the 6 parameters μ_f through symmetric polynomials

$$\sigma_k = \sum_{f_1 < \dots < f_k} \mu_{f_1} \dots \mu_{f_k} \quad (25)$$

As usual, $a_{ij} \equiv a_i - a_j$. Like the $SU(2)$ case, [4] the parameter $-\epsilon_1\epsilon_2$ should be absorbed into rescaling of Δ and w . Now one substitutes $w^2 = \Delta^3 - v^2$ with

$$\Delta = -\frac{(\alpha^2 + \beta^2)}{\epsilon_1\epsilon_2} \stackrel{(23)}{=} -\frac{1}{6\kappa^2\epsilon_1\epsilon_2} (a_1^2 + a_2^2 + a_3^2) \quad (26)$$

and

$$v^2 \stackrel{(9)}{=} -\frac{(\beta(\beta^2 - 3\alpha^2))^2}{(\epsilon_1\epsilon_2)^3} \stackrel{(23)}{=} -\frac{1}{108\kappa^6(\epsilon_1\epsilon_2)^3} (a_1 - a_2)^2 (a_2 - a_3)^2 (a_3 - a_1)^2 \quad (27)$$

and obtains for (21)

$$(a_1^2 + a_2^2 + a_3^2) \left(\tilde{p}(a_1^2 + a_2^2 + a_3^2)^3 - \tilde{s}a_{12}^2 a_{23}^2 a_{31}^2 \right) = a_{23}^2 P_6(a_1) + a_{13}^2 P_6(a_2) + a_{23}^2 P_6(a_3) \quad (28)$$

where $\tilde{p} = \frac{p+s}{12\kappa^2}$, $\tilde{s} = \frac{s}{6\kappa^2}$, and one should also impose condition (19), i.e. substitute $a_3 = -a_1 - a_2$. When all the external dimensions are vanishing (we still keep $\epsilon = 0$), the μ -parameters at the r.h.s. are vanishing as well:

$$\vec{\alpha}_1 = \dots = \vec{\alpha}_4 = \epsilon = 0 \implies \mu_1 = \dots = \mu_6 = 0 \quad (29)$$

Then the difference between the l.h.s. and the r.h.s. is a polynomial of degree 8 in a_1 and a_2 , equal to

$$2a_1^2 a_2^2 \left((80\tilde{p} - 13\tilde{s} - 4)(a_1^4 + a_2^4) + (128\tilde{p} + 17\tilde{s} + 2)a_1 a_2 (a_1^2 + a_2^2) + (152\tilde{p} + 32\tilde{s} + 5)a_1^2 a_2^2 \right) \quad (30)$$

For the AGT relation to be true this expression should vanish identically in a_1 and a_2 , i.e. the three coefficients should be made vanishing by adjusting just the two parameters $\tilde{p}$ and $\tilde{s}$. Surprisingly or not, the three equations are indeed consistent, and their common solution is $\tilde{p} = 1/72$, $\tilde{s} = -2/9$. This corresponds to $s/p = -8/9$, i.e. one obtains (24) in the form of the algebraic identity

$$\boxed{B^{(1)} = \frac{3\kappa^2}{2} \cdot \frac{\Delta(\Delta^3 - \frac{8}{9}w^2)}{\Delta^3 - w^2} \stackrel{(23)}{=} -\frac{1}{\epsilon_1\epsilon_2} \left(\frac{a_1^6}{a_{12}^2 a_{23}^2} + \frac{a_2^6}{a_{21}^2 a_{23}^2} + \frac{a_3^6}{a_{31}^2 a_{32}^2} \right) = Z^{(1)}} \quad (31)$$

The conformal block at the l.h.s. can be of course calculated from representation theory of the W_3 algebra, as we shall see in s.3 below.

2.4.3 Numerator. The case of arbitrary $\vec{\alpha}_2$ and $\vec{\alpha}_4$ with no ambiguity in the W -conformal block

If one switches on dimensions, the l.h.s. of (24) gets more complicated and additional contributions should be compensated by adjustment of six μ 's at the r.h.s. According to (10) and (11) with $q = 9/2$, the conformal block at the l.h.s. is now given by

$$2(\Delta_{\vec{\alpha}}^3 - w_{\vec{\alpha}}^2)B^{(1)} = \Delta^2 \bar{\Gamma}_{\vec{\alpha}_1, \vec{\alpha}_2; \vec{\alpha}}(L_{-1}) \Gamma_{\vec{\alpha}_3, \vec{\alpha}_4; \vec{\alpha}}(L_{-1}) - \\ - \frac{2w_{\vec{\alpha}}}{3} \left(\bar{\Gamma}_{\vec{\alpha}_1, \vec{\alpha}_2; \vec{\alpha}}(L_{-1}) \Gamma_{\vec{\alpha}_3, \vec{\alpha}_4; \vec{\alpha}}(W_{-1}) + \bar{\Gamma}_{\vec{\alpha}_1, \vec{\alpha}_2; \vec{\alpha}}(W_{-1}) \Gamma_{\vec{\alpha}_3, \vec{\alpha}_4; \vec{\alpha}}(L_{-1}) \right) + \frac{4\Delta_{\vec{\alpha}}}{9} \bar{\Gamma}_{\vec{\alpha}_1, \vec{\alpha}_2; \vec{\alpha}}(W_{-1}) \Gamma_{\vec{\alpha}_3, \vec{\alpha}_4; \vec{\alpha}}(W_{-1}) \quad (32)$$

where

$$\bar{\Gamma}_{\vec{\alpha}_1, \vec{\alpha}_2; \vec{\alpha}}(L_{-1}) = \Delta_{\vec{\alpha}} + \Delta_{\vec{\alpha}_1} - \Delta_{\vec{\alpha}_2} \quad (33)$$

and $\Gamma_{\vec{\alpha}_3, \vec{\alpha}_4; \vec{\alpha}}(L_{-1})$ is given by exactly the same formula, while $\Gamma(W_{-1})$ and $\bar{\Gamma}(W_{-1})$ require additional restrictions to be uniquely determined as was explained in the Introduction.

We begin with the case of $\vec{\alpha}_1 = \vec{\alpha}_3 = \vec{0}$, when the ambiguity is known not to show up. Then

$$\Gamma_{\vec{0}, \vec{\alpha}_4; \vec{\alpha}}(W_{-1}) = \xi w_{\vec{\alpha}} + \eta w_{\vec{\alpha}_2}, \quad \bar{\Gamma}_{\vec{0}, \vec{\alpha}_2; \vec{\alpha}}(W_{-1}) = \bar{\xi} w_{\vec{\alpha}} + \bar{\eta} w_{\vec{\alpha}_2} \quad (34)$$

with some coefficients $\xi, \bar{\xi}, \eta$ and $\bar{\eta}$. In order to find ξ , one can also put $\vec{\alpha}_2 = \vec{\alpha}_4 = 0$, then (32) turns into

$$B^{(1)} = \frac{\Delta}{2} \cdot \frac{\Delta^3 + w^2 \left(\left(\frac{2\xi}{3} - 1 \right) \left(\frac{2\bar{\xi}}{3} - 1 \right) - 1 \right)}{\Delta^3 - w^2} \quad (35)$$

Comparison with (31) implies that $\boxed{3\kappa^2 = 1}$ and $\boxed{\left(\frac{2\xi}{3} - 1 \right) \left(\frac{2\bar{\xi}}{3} - 1 \right) = 1/9}$.

Thus, we see that the AGT relation together with the basic definition (10) can only partly fix the coefficients in formulas for the conformal block at level one. Therefore, we glance at s.3.2 and fix the values $\xi = \bar{\xi}$. Then, $\xi = 1$ or 2 . The formulas of s.3.2 fix $\xi = 1$. Then, under switching on $\vec{\alpha}_2$ and $\vec{\alpha}_4$, (35) becomes

$$B^{(1)} = \frac{9\Delta^2(\Delta - \Delta_2)(\Delta - \Delta_4) - 6w(w + \eta w_2)(\Delta - \Delta_4) - 6w(w + \eta w_4)(\Delta - \Delta_2) + 4\Delta(w + \eta w_2)(w + \eta w_4)}{18(\Delta^3 - w^2)}$$

which should be now compared with the r.h.s. of (24). The resulting equations are identically satisfied for any a , provided

$$\begin{aligned} \sigma_1 &= 0, \\ \sigma_2 &= -(\alpha_2^2 + \beta_2^2 + \alpha_4^2 + \beta_4^2), \\ \sigma_3 &= -\frac{2}{3\sqrt{3}} \left(\bar{\eta} \alpha_2 (\alpha_2^2 - 3\beta_2^2) + \eta \alpha_4 (\alpha_4^2 - 3\beta_4^2) \right), \\ \sigma_4 &= (\alpha_2^2 + \beta_2^2)(\alpha_4^2 + \beta_4^2), \\ \sigma_5 &= \frac{2}{3\sqrt{3}} \left(\eta (\alpha_2^2 + \beta_2^2) \alpha_4 (\alpha_4^2 - 3\beta_4^2) + \bar{\eta} (\alpha_4^2 + \beta_4^2) \alpha_2 (\alpha_2^2 - 3\beta_2^2) \right), \\ \sigma_6 &= \frac{4\eta \bar{\eta}}{27} \alpha_2 (\alpha_2^2 - 3\beta_2^2) \alpha_4 (\alpha_4^2 - 3\beta_4^2) \end{aligned} \quad (36)$$

However, we still need to decompose these σ 's in μ according to (25). The AGT relation requires μ 's to be linear functions of $\vec{\alpha}$'s, and it is clear from the simple formula for σ_6 that this is indeed the case, with

$$\begin{aligned} \mu_1 &= -\bar{\eta} \frac{2}{\sqrt{3}} \alpha_2, \\ \mu_2 &= \bar{\eta} \frac{\alpha_2 + \beta_2 \sqrt{3}}{\sqrt{3}}, \\ \mu_3 &= \bar{\eta} \frac{\alpha_2 - \beta_2 \sqrt{3}}{\sqrt{3}}, \\ \mu_4 &= -\eta \frac{2}{\sqrt{3}} \alpha_4, \\ \mu_5 &= \eta \frac{\alpha_4 + \beta_4 \sqrt{3}}{\sqrt{3}}, \\ \mu_6 &= \eta \frac{\alpha_4 - \beta_4 \sqrt{3}}{\sqrt{3}} \end{aligned} \quad (37)$$

and $\boxed{\eta, \bar{\eta} = \pm 1}$. This constraint is not a surprise since in (36) η and $\bar{\eta}$ appear rather irregularly. This solution also reproduces all other σ 's and it is unique up to the sign choice for $\eta, \bar{\eta}$ and up to 6! permutations of μ 's.

2.4.4 The case of $\xi = 2$

Let us now see what happens if one chooses $\xi = \bar{\xi} = 2$ in (34). It turns out that this time the overdefined system of equations for six σ 's have *no* solution. Solution appears if one takes into account an extra factor

$$Z^{U(1)} = (1 - x)^{-\nu} \quad (38)$$

in the AGT relation,

$$\mathcal{B} = Z^{U(1)} Z^{SU(N)} \quad (39)$$

At level one introduction of ν implies that (24) is deformed into

$$B^{(1)} = Z^{(1)} + \nu \quad (40)$$

If $\nu \neq 0$ is allowed, then $\xi = 2$ also provides a solution to (40), actually with the same parameters μ as in (37) and with

$$\nu = \alpha_2^2 + \alpha_4^2 + \beta_2^2 + \beta_4^2 = \Delta_2 + \Delta_4 \quad (41)$$

The origin for this other solution is that it corresponds to the other choice of the special states, 2 and 4. This is why this value of ξ is not consistent with the table (111) where the special states are chosen to be 1 and 3.

2.4.5 The $U(1)$ factor ν and projective transformations

Thus we observe the same phenomenon as in the $SU(2)$ case in [4]: if one makes the simultaneous interchange of external lines $\vec{\alpha}_1 \leftrightarrow \vec{\alpha}_2$ and $\vec{\alpha}_3 \leftrightarrow \vec{\alpha}_4$ a simple ν (zero in the particular case of $\vec{\alpha}_1 = \vec{\alpha}_3 = 0$ which we are now considering) turns into a sophisticated one. In principle this is nothing but a result of modular transformation of the conformal block. Indeed, the generic four-point function is [7, 10]

$$\langle V_{\Delta_1}(z_1) V_{\Delta_2}(z_2) V_{\Delta_3}(z_3) V_{\Delta_4}(z_4) \rangle = \prod_{i < j}^4 (z_i - z_j)^{d_{ij}} (\bar{z}_i - \bar{z}_j)^{\bar{d}_{ij}} G(x, \bar{x}) \quad (42)$$

where $d_{34} = d_{13} = 0$, $d_{14} = -2\Delta_2$, $d_{24} = \Delta_1 + \Delta_3 - \Delta_2 - \Delta_4$, $d_{34} = \Delta_1 + \Delta_2 - \Delta_3 - \Delta_4$, $d_{23} = \Delta_4 - \Delta_2 - \Delta_1 - \Delta_3$, $x = \frac{(z_1 - z_2)(z_3 - z_4)}{(z_3 - z_2)(z_1 - z_4)}$ and similarly for the complex conjugated part. $G(x, \bar{x})$ is a bilinear combination of the conformal blocks $B(\{\Delta_i\}, \Delta, c; x)$ and $B(\{\bar{\Delta}_i\}, \bar{\Delta}, c; \bar{x})$. Due to the projective invariance, one can choose three of these four points arbitrarily. If choosing $z_4 = \infty$, $z_3 = 1$ and $z_2 = 0$, one obtains $z_1 = x$ and the 4-point correlator (42) becomes

$$(1 - x)^{d_{12}} \bar{x}^{\bar{d}_{13}} (1 - \bar{x})^{\bar{d}_{12}} x^{d_{13}} G(x, \bar{x}) \quad (43)$$

The simultaneous interchange of external lines $\vec{\alpha}_1 \leftrightarrow \vec{\alpha}_2$ and $\vec{\alpha}_3 \leftrightarrow \vec{\alpha}_4$ in formula (42) does not change x and $\bar{x}$ and leads to the factor of $(1 - x)^{d_{24}}$ instead of $(1 - x)^{d_{13}}$ ($d_{13} = 0$) which is exactly $\nu = -d_{24}$ in (41) (since $\Delta_1 = \Delta_3 = 0$ there).

2.4.6 General case, all $\vec{\alpha} \neq 0$, $\vec{\alpha}_1$ and $\vec{\alpha}_3$ special

We now come to the general case of all external $\vec{\alpha}$ switched on. In this case, one needs some ansatz for the 3-point functions $\Gamma(W_{-1})$, $\bar{\Gamma}(W_{-1})$. However, the number of parameters is large enough and the adjusting procedure fails to be very effective to define the coefficients. As we know from the $SU(2)$ case, restrictions get very strong when one includes the AGT relation at higher levels [4], see s.5.

Therefore, we return to a more straightforward approach and just take the true value of the 3-point function from the conformal field theory analysis, see [5] and s.3.2 below. Thus, we take the vertices from s.3.2 and fix the special states to have $\beta_{1,3} = r_{1,3} \alpha_{1,3}$ with some fixed $r_{1,3}$.




Figure 1: Roots and minimal weights of $sl(3)$. The roots have the length $\sqrt{2}$, the minimal weights have $\sqrt{2/3}$. All angles are integer multiples of 30° . The two simple roots and the corresponding two fundamental weights are marked by black circles. The third positive root, marked with a white circle is also $\vec{\rho}$, the half-sum of all (three) positive roots or the sum of (two) fundamental weights. The vectors $\vec{\lambda}_{1,2,3}$ are shown by thick lines, they form a Weyl-invariant triple of minimal weights.

Then AGT relation is satisfied, provided there is a certain linear relation between the sextuplets of parameters $\alpha_1, \vec{\alpha}_2, \alpha_3, \vec{\alpha}_4$ and $\mu_1, \dots, \mu_6$. One of the possible solutions is

$$\begin{aligned}
\mu_1 &= \frac{2}{\sqrt{3}}(\alpha_2 - \alpha_1), \\
\mu_2 &= -\frac{\alpha_2 + 2\alpha_1 + \beta_2\sqrt{3}}{\sqrt{3}}, \\
\mu_3 &= -\frac{\alpha_2 + 2\alpha_1 - \beta_2\sqrt{3}}{\sqrt{3}}, \\
\mu_4 &= -\frac{2}{\sqrt{3}}(\alpha_4 + \alpha_3), \\
\mu_5 &= \frac{\alpha_4 + 2\alpha_3 + \beta_4\sqrt{3}}{\sqrt{3}}, \\
\mu_6 &= \frac{\alpha_4 + 2\alpha_3 - \beta_4\sqrt{3}}{\sqrt{3}}
\end{aligned} \tag{44}$$

In this solution, the $U(1)$ -factor shows up in the level one AGT relation (40) with $\nu = -4\alpha_1\alpha_3$, while

$$r_1 = -\sqrt{3}, \quad r_3 = \sqrt{3} \tag{45}$$

Let us introduce the three vectors, see Fig.1

$$\vec{\lambda}_1 = \frac{1}{\sqrt{6}}(1, \sqrt{3}), \quad \vec{\lambda}_2 = \frac{1}{\sqrt{6}}(1, -\sqrt{3}), \quad \vec{\lambda}_3 = \sqrt{\frac{2}{3}}(-1, 0) \tag{46}$$

They are three (out of six) minimal vectors in the weight lattice of $sl(3)$. Then a_i in (23) with $\kappa = 1/\sqrt{3}$ and μ_f in (44) can be represented as scalar products,

$$a_{1,2,3} = \sqrt{2}\vec{\lambda}_{1,2,3}\vec{\alpha} = (\sqrt{2}\vec{\alpha})\vec{\lambda}_{1,2,3} \tag{47}$$

and

$$\begin{aligned}
\mu_{1,2,3} &= -(\sqrt{2}\vec{\alpha}_2)\vec{\lambda}_{1,2,3} - \frac{2}{\sqrt{3}}\alpha_1, \\
\mu_{4,5,6} &= (\sqrt{2}\vec{\alpha}_4)\vec{\lambda}_{1,2,3} - \frac{2}{\sqrt{3}}\alpha_3
\end{aligned} \tag{48}$$

Let us remind that the primary exponentials are $V_{\vec{\alpha}} = e^{(\sqrt{2}\vec{\alpha})\phi}$, this can make these formulas looking more natural. The eigenvalues of the $W^{(3)}$ operator are

$$w_{\vec{\alpha}} = -3\sqrt{6}(\vec{\alpha}\vec{\lambda}_1)(\vec{\alpha}\vec{\lambda}_2)(\vec{\alpha}\vec{\lambda}_3) = \alpha(\alpha^2 - 3\beta^2) \tag{49}$$

Note that one could choose the pair of the *special* states in a way, different from (45), that is, $r_{1,3} = 0, \pm\sqrt{3}$. We return to complete analysis in s.4.2 below, and the full final result is summarized in the Conclusion, eqs.(131)-(134).

2.5 Intermediate summary

This ends the explicit check of the AGT conjecture for the 4-point conformal block of W_3 -algebra with $c = 2$ at level one: it is indeed expressed through Nekrasov's special functions with $\vec{a}$ and μ_f linearly expressed through $\vec{a}$'s. Technical generalizations can go in four obvious directions.

First, one can check the AGT relation between the structure constants of conformal model and "perturbative" Nekrasov's functions. This relation refers to the 3-point functions of particular conformal model: Liouville in the $SU(2)$ and conformal Toda in the more general $SU(N)$ case, which are not yet known in full generality. It was also discussed in some detail in [2].

The second and third, one can either switch on $\epsilon \neq 0$ or proceed to higher levels or do the both, and repeat the calculations made for the $SU(2)$ case in [4]. Fourth, one can consider the $SU(N)/W_N$ examples with $N > 3$. These are straightforward exercises in representation theory.

In order proceed to a discussion of the second and third directions in ss.4 and 5 respectively, we need some additional information from the theory of W -algebras, it is reminded in [5] and, more briefly, in the following section 3. As to the fourth direction, we discuss only one, but very interesting subject in s.6: it concerns a general role of the Nekrasov functions and, more generally, matrix-model τ -functions as a modern substitute for hypergeometric functions. As a particular application, s.6 comments on the complete proof [6] of the AGT relation for the special case when the conformal blocks are hypergeometric functions (rather than generic "hypergeometric integrals" [26, 27], to which they are supposedly related in the Dotsenko-Fateev approach).

3 Some CFT considerations

In s.2 we constructed conformal blocks at level one mostly from dimensional considerations and demonstrated that remaining few parameters can be adjusted so that the AGT relations are fulfilled. Moreover, even at level one restrictions are rather strong, they actually become exhaustive already at level two. We did this on purpose: to demonstrate that the AGT relations can in the future become a powerful alternative approach to conformal field theory problems, which are pretty hard when attacked directly. Still, conformal blocks are conformal blocks and they can of course be evaluated by the standard CFT methods. In this section we briefly sketch how to evaluate the relevant quantities: the 3-point functions Γ and $\bar{\Gamma}$ at level one and the Shapovalov matrix. The details can be found in [5]. For original papers and reviews see [7, 10, 12, 14, 16, 17].

Note that we use somewhat non-standard normalizations, adjusted to maximally simplify the W -algebra formulas. The price for this is certain deviation from conventions in some other papers. We illustrate our conventions with the example of free field theory in the Appendix.

3.1 W -algebra and Shapovalov form

In s.2.2 we defined the parameter q in (11) from the requirement that the corresponding Kac determinant decomposes nicely with the help of (9). In CFT q is calculated from the structure constants of W_3 -algebra. However, the simple ones (12) are not sufficient, one needs also the remaining, non-linear commutation relation [13]-[17]:

$$\begin{aligned} \frac{2}{9} [W_n, W_m] = (n-m) \left\{ \frac{16}{22+5c} \Lambda_{n+m} + \left(\frac{(n+m+2)(n+m+3)}{15} - \frac{(n+2)(m+2)}{6} \right) L_{n+m} \right\} + \\ + \frac{c}{3 \cdot 5!} n(n^2-1)(n^2-4) \delta_{n+m,0} \end{aligned} \quad (50)$$

Here Λ_n is a remnant of the $W^{(4)}$ operator. It does not exist as an independent operator in the W_3 algebra for the same reason that $\text{tr } J^4 = \frac{1}{2} (\text{tr } J^2)^2$ for a 3×3 traceless matrix J . In fact,

$$\Lambda_n = \sum_{k=-\infty}^{\infty} : L_k L_{n-k} : + \frac{x_n}{5} L_n, \quad (51)$$

with $x_{2l} = 1-l^2$, $x_{2l+1} = (2+l)(1-l)$. We use the notation from [17], but our normalization of the W -operator is different by a factor of $3/\sqrt{2}$. Note for $n = 0$ that the term with the normal ordering contains an item L_0^2 which *does* contribute to the vacuum average:

$$\langle V_\alpha \Lambda_0 V_\alpha \rangle = \Delta_\alpha^2 + \frac{1}{5} \Delta_\alpha = \Delta_\alpha \left(\Delta_\alpha + \frac{1}{5} \right) \quad (52)$$

Thus one obtains for the "difficult" element of the Shapovalov matrix (11), (see [4] where a similar calculation for the Virasoro group is reminded in detail):

$$\langle W_{-1} V_{\bar{\alpha}} | W_{-1} V_{\bar{\alpha}} \rangle = \langle V_{\bar{\alpha}} | [W_1, W_{-1}] V_{\bar{\alpha}} \rangle = \frac{9\Delta_{\bar{\alpha}}}{2} \left(\frac{32}{22+5c} \left(\Delta_{\bar{\alpha}} + \frac{1}{5} \right) - \frac{1}{5} \right) \equiv \frac{9D_{\bar{\alpha}}\Delta_{\bar{\alpha}}}{2} \xrightarrow{c=2} \frac{9}{2}\Delta_{\bar{\alpha}}^2 \quad (53)$$

in accordance with the *guess* in s.2.2. The other three entries of the table (11) do *not* depend on c . Likewise, more generally,

$$\langle W_{-n} V_{\bar{\alpha}} | W_{-n} V_{\bar{\alpha}} \rangle = \frac{9n\Delta_{\bar{\alpha}}}{2} \left(\frac{32}{22+5c} \left(\Delta_{\bar{\alpha}} + \frac{1}{5} \right) + \frac{5n^2-8}{15} \right) = \frac{9D_{\bar{\alpha}}\Delta_{\bar{\alpha}}}{2} = \frac{9n\Delta_{\bar{\alpha}}}{2} \left(D_{\bar{\alpha}} + \frac{n^2-1}{3} \right) \quad (54)$$

We introduced here a peculiar parameter

$$D_{\bar{\alpha}} \equiv \frac{32}{22+5c} \left(\Delta_{\bar{\alpha}} + \frac{1}{5} \right) - \frac{1}{5} \quad (55)$$

which helps to get some formulas at low levels shorter (however, D is nothing like a universal effective variable to absorb, say, the c -dependence; at most, it is a convenient abbreviation for limited purposes). Two other expressions of this kind are:

$$\begin{aligned} [W_0, W_{-1}] V_{\bar{\alpha}} &= \frac{9D}{2} L_{-1} V_{\bar{\alpha}}, \\ W_1 W_{-1} V_{\bar{\alpha}} &= [W_1, W_{-1}] V_{\bar{\alpha}} = \frac{9D\Delta}{2} V_{\bar{\alpha}} \end{aligned} \quad (56)$$

where $V_{\bar{\alpha}}$ is arbitrary primary.

3.2 3-point functions at level one

Three point functions of interest for us are evaluated simply by moving integration contours. In (10) we need the two *types* of such functions, Γ and $\bar{\Gamma}$, which are different in the W -sector, see [5].

We begin with the $\Gamma(L_1)$, i.e. with the *stress-tensor* insertion into a *correlator* of three primary fields (we write it for the operators V_3 and V_4 , rather than V_1 and V_2 , since this is the form in which we actually need it in (10), also location of the Virasoro operator is underlined to make the formula readable):

$$\begin{aligned} \langle \underline{L_n V_{\alpha}(0)} V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle &= \oint_0 x^{n+1} dx \langle T(x) V_{\alpha}(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle = \\ &= - \left(\oint_1 x^{n+1} dx + \oint_{\infty} x^{n+1} dx \right) \langle T(x) V_{\alpha}(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle = \\ &= - \sum_k \oint_1 \frac{x^{n+1} dx}{(x-1)^{k+2}} \langle V_{\alpha}(0) \underline{L_k V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle + \sum_k \oint_{\infty} x^{k-2} x^{n+1} dx \langle V_{\alpha}(0) V_{\alpha_3}(1) \underline{L_k V_{\alpha_4}(\infty)} \rangle = \\ &= -(n+1) \Delta_3 \langle V_{\alpha}(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle - \langle V_{\alpha}(0) \underline{L_{-1} V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle + \langle V_{\alpha}(0) V_{\alpha_3}(1) \underline{L_{-n} V_{\alpha_4}(\infty)} \rangle \end{aligned} \quad (57)$$

where we used the fact that primaries are annihilated by L_n with $n > 0$ and the action of L_0 produces the dimension: because of this, the first sum in the third line is reduced to two terms. Only one term contributes to the second sum even without these conditions. Now one can substitute $n = -1$, $n = 0$ and $n = 1$ to get (in an obvious abbreviated notation)

$$\begin{aligned} \langle L_{-1} V_{\alpha}(0) \rangle &= -\langle L_{-1} V_{\alpha_3}(1) \rangle, \\ \Delta &= -\Delta_3 - \langle L_{-1} V_{\alpha_3}(1) \rangle + \Delta_4, \\ 0 &= -2\Delta_3 - \langle L_{-1} V_{\alpha_3}(1) \rangle + \langle L_{-1} V_{\alpha_4}(\infty) \rangle \end{aligned} \quad (58)$$

which can be resolved for any of the three positions of the L_{-1} operator:

$$\begin{aligned} \langle \underline{L_{-1} V_{\alpha}(0)} V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle &= (\Delta + \Delta_3 - \Delta_4) \langle V_{\alpha}(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle, \\ \langle V_{\alpha}(0) \underline{L_{-1} V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle &= (-\Delta - \Delta_3 + \Delta_4) \langle V_{\alpha}(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle, \\ \langle V_{\alpha}(0) V_{\alpha_3}(1) \underline{L_{-1} V_{\alpha_4}(\infty)} \rangle &= (-\Delta + \Delta_3 + \Delta_4) \langle V_{\alpha}(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle \end{aligned} \quad (59)$$

It is the first position that we denote through $\Gamma_{\alpha_3, \alpha_4; \alpha}(L_{-1})$ in the main text, and it is the second (boxed) one that we use when constructing $\Gamma_{\alpha_3, \tilde{\alpha}_4; \tilde{\alpha}}(W_{-1})$ for the special state $\tilde{\alpha}_3$.

Similarly, the $\Gamma(W_{-1})$ -function results from the study of W -operator insertions:

$$\begin{aligned}
\langle \underline{W_n V_\alpha(0)} V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle &= \oint_0 x^{n+2} dx \langle W(x) V_\alpha(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle = \\
&= - \left(\oint_1 x^{n+2} dx + \oint_\infty x^{n+2} dx \right) \langle W(x) V_\alpha(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle = \\
&= - \sum_k \oint_1 \frac{x^{n+2} dx}{(x-1)^{k+3}} \langle V_\alpha(0) \underline{W_k V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle \boxed{-} \sum_k \oint_\infty x^{k-3} x^{n+2} dx \langle V_\alpha(0) V_{\alpha_3}(1) \underline{W_k V_{\alpha_4}(\infty)} \rangle = \\
&= - \frac{(n+1)(n+2)w_3}{2} \langle V_\alpha(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle - (n+2) \langle V_\alpha(0) \underline{W_{-1} V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle - \\
&\quad - \langle V_\alpha(0) \underline{W_{-2} V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle - \langle V_\alpha(0) V_{\alpha_3}(1) \underline{W_{-n} V_{\alpha_4}(\infty)} \rangle \quad (60)
\end{aligned}$$

Note that the sign in the box is *different* from the Virasoro case, because the odd-spin 3-differential $W(z)$ is transformed differently from the even-spin 2-differential $T(z)$ under the change $z \rightarrow 1/z$, $dz/z \rightarrow -dz/z$. Applying the same trick, i.e. putting $n = -1, 0, 1$ one gets in abbreviated notation:

$$\begin{aligned}
\langle W_{-1} V_\alpha(0) \rangle &= & -\langle W_{-1} V_{\alpha_3}(1) \rangle & -\langle W_{-2} V_{\alpha_3}(1) \rangle, \\
w &= & -w_3 & -2\langle W_{-1} V_{\alpha_3}(1) \rangle & -\langle W_{-2} V_{\alpha_3}(1) \rangle & -w_4, \\
0 &= & -3w_3 & -3\langle W_{-1} V_{\alpha_3}(1) \rangle & -\langle W_{-2} V_{\alpha_3}(1) \rangle & -\langle W_{-1} V_{\alpha_4}(\infty) \rangle
\end{aligned} \quad (61)$$

This time, however, the four unknowns can not be found from these three equations: we can only exclude terms with W_{-2} obtain pair relations between the terms with W_{-1} :

$$\begin{aligned}
\boxed{\langle \underline{W_{-1} V_\alpha(0)} V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle} &= \left(w + w_3 + w_4 \right) \langle V_\alpha(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle + \langle V_\alpha(0) \underline{W_{-1} V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle, \\
\langle \underline{W_{-1} V_\alpha(0)} V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle &= \left(2w - w_3 + 2w_4 \right) \langle V_\alpha(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle - \langle V_\alpha(0) V_{\alpha_3}(1) \underline{W_{-1} V_{\alpha_4}(\infty)} \rangle, \\
\langle V_\alpha(0) \underline{W_{-1} V_{\alpha_3}(1)} V_{\alpha_4}(\infty) \rangle &= \left(w - 2w_3 + w_4 \right) \langle V_\alpha(0) V_{\alpha_3}(1) V_{\alpha_4}(\infty) \rangle - \langle V_\alpha(0) V_{\alpha_3}(1) \underline{W_{-1} V_{\alpha_4}(\infty)} \rangle \quad (62)
\end{aligned}$$

Only if an additional constraint like (4) is imposed on any *one* of the three states, all correlators can be unambiguously defined in a universal way: through Δ 's and w 's (or, what is equivalent, through the parameters $\tilde{\alpha}$). It is the first (boxed) of these formulas that was relevant for our consideration in s.2.4.

The triple vertices of another type $\bar{\Gamma}$ are matrix elements rather than correlators (averages), they also obey a Ward identity, similar to (57), but not exactly the same. We refer to [5] for details of the derivation and present here only the answers, which we need the most:

$$\begin{aligned}
\boxed{\bar{\Gamma}(L_{-1})_{\alpha_1 \alpha_2; \alpha} = \langle \underline{L_{-1} V_\alpha} | V_{\alpha_1}(1) V_{\alpha_2}(0) \rangle} &= \left(\Delta_\alpha + \Delta_1 - \Delta_2 \right) \langle V_\alpha | V_1(1) V_2(0) \rangle, \\
\langle V_\alpha | \underline{(L_{-1} V_1)(1)} V_2(0) \rangle &= \left(\Delta_\alpha - \Delta_1 - \Delta_2 \right) \langle V_\alpha | V_1(1) V_2(0) \rangle, \quad (63)
\end{aligned}$$

$$\boxed{\bar{\Gamma}(W_{-1})_{\alpha_1 \alpha_2; \alpha} = \langle \underline{W_{-1} V_\alpha} | V_{\alpha_1}(1) V_{\alpha_2}(0) \rangle} = (w_\alpha + 2w_1 - w_2) \langle V_\alpha | V_1(1) V_2(0) \rangle + \langle V_\alpha | \underline{(W_{-1} V_1)(1)} V_2(0) \rangle$$

The second formula in (63) is used to handle the situation when V_{α_1} is *special*.

3.3 Shapovalov matrix at level two

Similar relations at level two are more sophisticated, but can be straightforwardly derived in the same way. They are listed in a separate paper [5], where they are also checked with the help of the free field model. This check is of importance to make because the derivation is rather tedious and mistakes are not so easy to exclude. We present and use these partly-validated answers in s.5 below.

It remains to evaluate the W_3 Shapovalov form. At level one it is very simple: if expressed in terms of D from eq.(55), it turns from (11) into

$$\begin{array}{|c|c|} \hline 2\Delta & 3w \\ \hline 3w & 9D\Delta/2 \\ \hline \end{array} \quad (64)$$

for all values of c . At level two there are five "W₃-Young diagrams" $\mathcal{Y}$ with $|\mathcal{Y}| = 2$ and one gets a 5×5 matrix:

$Q_\alpha(\mathcal{Y}, \mathcal{Y}')$	$L_{-2}V_\alpha$	$L_{-1}^2V_\alpha$	$L_{-1}W_{-1}V_\alpha$	$W_{-2}V_\alpha$	$W_{-1}^2V_\alpha$
$L_{-2}V_\alpha$	$4\Delta + \frac{c}{2}$	6Δ	$9w$	$6w$	$\frac{45D\Delta}{2}$
$L_{-1}^2V_\alpha$	6Δ	$4\Delta(2\Delta + 1)$	$6w(2\Delta + 1)$	$12w$	$9(3D\Delta + 2w^2)$
$L_{-1}W_{-1}V_\alpha$	$9w$	$6w(2\Delta + 1)$	$9(D\Delta^2 + D\Delta + w^2)$	$18D\Delta$	$\frac{27Dw}{2}(2\Delta + 3)$
$W_{-2}V_\alpha$	$6w$	$12w$	$18D\Delta$	$9\Delta(D + 1)$	$\frac{27w}{2}(3D + 1)$
$W_{-1}^2V_\alpha$	$\frac{45D\Delta}{2}$	$9(3D\Delta + 2w^2)$	$\frac{27Dw}{2}(2\Delta + 3)$	$\frac{27w}{2}(3D + 1)$	$\frac{81}{4}D^2\Delta(2\Delta + 1) + \frac{81\kappa}{4}(D\Delta(\Delta + 1) + 4w^2)$

(65)

where special notation κ is introduced for a peculiar combination

$$\kappa = \frac{32}{22 + 5c} \xrightarrow{c=2} 1, \quad \text{then} \quad D = \kappa \left(\Delta + \frac{1}{5} \right) - \frac{1}{5} \xrightarrow{c=2} \Delta \quad (66)$$

We describe here only evaluation of the tricky entries of (65), which involves commutators of W -operators. The first such example is

$$< L_{-2}V_\alpha | W_{-1}^2V_\alpha > = < \alpha | L_2W_{-1}^2 | \alpha > \stackrel{(12)}{=} < \alpha | 5W_1W_{-1} | \alpha > \stackrel{(50)}{=} \frac{45}{2}D\Delta \quad (67)$$

At the last stage we could of course directly use (56). As in [4] we use obvious abbreviated notation, which we now abbreviate even further. Similarly,

$$L_1^2W_{-1}^2 \stackrel{(12)}{=} L_1(W_{-1}L_{-1} + 3W_0)W_{-1} = (L_1W_{-1})^2 + 3(L_1W_{-1})W_0 + 3L_1[W_0, W_{-1}] \quad (68)$$

The first two terms are straightforward: they give $(3w)^2 + 3(3w)w = 18w^2$ when acting on a primary field. The last term is trickier: according to (50), the commutator turns into $\frac{9}{2}(\kappa/2)\Lambda_{-1} - (1/5)L_{-1}$ and then into $\frac{9}{2}L_{-1}(\kappa(\Delta + 1/5) - 1/5) = \frac{9D}{2}L_{-1}$. Finally, $3L_1L_{-1} \rightarrow 6\Delta$ and one obtains for this entry of the table: $18w^2 + 27D\Delta$. Alternatively, in (68) one could push W_0 to the left: $3L_1W_0W_{-1} = 3(W_0L_1 + 2W_1)W_{-1} \rightarrow 9w^2 + 27D\Delta$, which reproduces the same result in a simpler way.

The other entries of the matrix (65) are calculated in a similar way:

$$W_1L_1L_{-1}W_{-1} = W_1(2L_0 + L_{-1}L_1)W_{-1} \rightarrow 2W_1(W_{-1}(L_0 + 1)) + 3W_1L_{-1}W_0 \rightarrow 9D\Delta(\Delta + 1) + 9w^2; \quad (69)$$

$$W_1L_1W_{-2} \rightarrow 4W_1W_{-1} \rightarrow 18D\Delta; \quad (70)$$

$$W_1L_1W_{-1}^2 \rightarrow W_1(W_{-1}L_1 + 3W_0)W_{-1} \rightarrow \frac{27}{2}wD\Delta + \frac{27}{2}wD\Delta + \frac{27D}{2}W_1L_{-1} \rightarrow \frac{27wD}{2}(2\Delta + 3) \quad (71)$$

$$W_1^2W_{-2} \rightarrow W_1[W_1, W_{-2}] = \frac{27}{2}W_1 \left(\frac{\kappa}{2}\Lambda_{-1} + \frac{2}{15}L_{-1} \right) \rightarrow \frac{27}{2}W_1L_{-1} \left(\kappa \left(\Delta + \frac{1}{5} \right) + \frac{2}{15} \right) \rightarrow \frac{81w}{2} \left(D + \frac{1}{3} \right) \quad (72)$$

$$W_1^2W_{-1}^2 = W_1 \left(W_{-1}W_1 + \frac{9}{2}(\kappa\Lambda_0 - \frac{1}{5}L_0) \right) W_{-1} \rightarrow \left(\frac{9D\Delta}{2} \right)^2 + \frac{9\kappa}{2}W_1\Lambda_0W_{-1} - \frac{9}{10} \cdot \frac{9}{2}D\Delta(\Delta + 1) \rightarrow \quad (73)$$

The middle term in the last expression is a little tedious: since Λ_0 acts on a first descendant rather than primary, one needs to include also a term from the normal ordered part: $W_1 \Lambda_0 W_{-1} \rightarrow W_1 (L_0^2 + \frac{1}{5} L_0 + 2L_{-1} L_1) W_{-1} \rightarrow \frac{9D\Delta}{2} (\Delta + \frac{6}{5}) (\Delta + 1) + 18w^2$. Collecting all terms, one finally gets

$$W_1^2 W_{-1}^2 \rightarrow \frac{81}{4} (D^2 \Delta (2\Delta + 1) + \kappa (D\Delta (\Delta + 1) + 4w^2)) \quad (74)$$

4 AGT relation to $c \neq 2$ at level one

As was already mentioned, going to arbitrary central charges c and to higher levels, i.e. to higher powers of x and bigger sizes of the generalized Young diagrams, is a straightforward, but tedious exercise in group theory. The crucial difficult fact is actually well known: the Kac determinant is *always* (i.e. for arbitrary level and for arbitrary W_N algebra) nicely factorized in terms of $\vec{\alpha}$ -variables, so that its zeroes are always given by an integer combination $\vec{\alpha} \vec{e}_i = m\epsilon_1 + n\epsilon_2$. This fact is remarkable, but well known, its best heuristic "explanation" comes from the theory of free fields [12], but algebraically it looks somewhat artificial.

4.1 No external lines, $\vec{\alpha}_1, \dots, \vec{\alpha}_4 = 0$

At level one for arbitrary c it follows from existence of the deformation of (22):

$$\Delta^3 - w^2 = (\alpha^2 + \beta^2)^3 - (\alpha(\alpha^2 - 3\beta^2))^2 = (\beta(\beta^2 - 3\alpha^2))^2 \longrightarrow$$

$$(\alpha^2 + \beta^2 - Q^2)^2 \left(\alpha^2 + \beta^2 - \frac{1}{4} Q^2 \right) - \alpha^2 (\alpha^2 - 3\beta^2)^2 = (\beta^2 - Q^2/4) \left((\beta - Q)^2 - 3\alpha^2 \right) \left((\beta + Q)^2 - 3\alpha^2 \right) \quad (75)$$

The l.h.s. of this deformed relation is interpreted in accordance with (64), as determinant of the Shapovalov matrix,

$$\begin{aligned} D\Delta^2 - w^2 &= \left(\frac{1}{1 - \frac{15}{4}Q^2} (\alpha^2 + \beta^2 - Q^2 + \frac{1}{5}) - \frac{1}{5} \right) (\alpha^2 + \beta^2 - Q^2)^2 - \frac{1}{1 - \frac{15}{4}Q^2} \alpha^2 (\alpha^2 - 3\beta^2)^2 = \\ &= \frac{1}{1 - \frac{15}{4}Q^2} \left\{ (\alpha^2 + \beta^2 - Q^2)^2 \left(\alpha^2 + \beta^2 - \frac{1}{4}Q^2 \right) - \alpha^2 (\alpha^2 - 3\beta^2)^2 \right\} \end{aligned} \quad (76)$$

i.e. $c = 2(1 - 12Q^2)$ and $\Delta = \alpha^2 + \beta^2 - Q^2$. Note that the eigenvalue w and, thus, the operator $W^{(3)}$ itself also acquire the c -dependent factor $\sqrt{\frac{32}{22+5c}}$.

In the AGT relation, the r.h.s. of (75) should match with a deformation of square of the Van-der-Monde determinant in the Nekrasov formula, given by the denominator in eq.(17): instead of (20) one now has

$$\begin{aligned} &(\beta^2 - Q^2/4) \left((\beta - Q)^2 - 3\alpha^2 \right) \left((\beta + Q)^2 - 3\alpha^2 \right) \sim \\ &\sim (a_{12} - \epsilon) a_{12} (a_{12} + \epsilon) (a_{23} - \epsilon) a_{23} (a_{23} + \epsilon) (a_{31} - \epsilon) a_{31} (a_{31} + \epsilon) \end{aligned} \quad (77)$$

Note that the number of different structures at the r.h.s. (77), i.e. in the denominator of the Nekrasov formula, increased from six to nine with switching on a non-zero ϵ . Some three zeroes of the denominator should be canceled by the numerator of the Nekrasov formula, and only the remaining six should match the six zeroes at the l.h.s. of (77). From experience with the $SU(2)$ case in [4], one can guess that *irrelevant* three factors at the r.h.s. of (77) are nothing but $a_{12} a_{23} a_{31} = \Delta(a)$, and the mapping (23) should be deformed so that

$$\begin{pmatrix} a_{12} = -2\beta \\ a_{23} = \beta + \alpha\sqrt{3} \\ a_{31} = \beta - \alpha\sqrt{3} \end{pmatrix} \longrightarrow \begin{pmatrix} a_{12} + \epsilon = -2\beta + Q \\ a_{23} + \epsilon = \beta + Q + \alpha\sqrt{3} \\ a_{31} + \epsilon = \beta + Q - \alpha\sqrt{3} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} a_{12} - \epsilon = -2\beta - Q \\ a_{23} - \epsilon = \beta - Q + \alpha\sqrt{3} \\ a_{31} - \epsilon = \beta - Q - \alpha\sqrt{3} \end{pmatrix} \quad (78)$$

i.e. $\epsilon = Q$. In fact, after appropriate rescalings, see [4],

$$Q^2 = \frac{\epsilon^2}{-\epsilon_1 \epsilon_2} \quad (79)$$

and

$$\begin{aligned} \Delta_{\vec{\alpha}} &= \frac{\alpha^2 + \beta^2 - Q^2}{-\epsilon_1 \epsilon_2}, \\ w_{\vec{\alpha}} &= \sqrt{\frac{\kappa}{-\epsilon_1 \epsilon_2}} \alpha (\alpha^2 - 3\beta^2) \end{aligned} \quad (80)$$

Note that (23) actually remains intact:

$$\begin{aligned} a_1 &= \kappa(\alpha - \beta\sqrt{3}), \\ a_2 &= \kappa(\alpha + \beta\sqrt{3}), \\ a_3 &= -2\kappa\alpha \end{aligned} \tag{81}$$

with $\kappa = 1/\sqrt{3}$.

For generic c the AGT relation (24) turns into

$$\begin{aligned} & \frac{1}{2} \cdot \frac{\Delta(D\Delta^2 - \frac{8}{9}w^2)}{D\Delta^2 - w^2} = \\ &= \frac{1}{\epsilon_1\epsilon_2} \left(\frac{P_6(a_1)}{a_{12}(a_{12} + \epsilon)a_{13}(a_{13} + \epsilon)} + \frac{P_6(a_2)}{a_{21}(a_{21} + \epsilon)a_{23}(a_{23} + \epsilon)} + \frac{P_6(a_3)}{a_{31}(a_{31} + \epsilon)a_{32}(a_{32} + \epsilon)} \right) + \nu \end{aligned} \tag{82}$$

where the conformal block at the l.h.s. is the deformation of (31) and the r.h.s. is the level one Nekrasov function for non-vanishing $\epsilon = \epsilon_1 + \epsilon_2$. The matching of denominators is guaranteed by the basic relation (75), to which we adjusted all our choices of dimensions and the AGT relation $\vec{a} \xleftrightarrow{(81)} \vec{\alpha}$. Therefore (82) defines the polynomial $P(a)$ and the $U(1)$ parameter ν . Like in the $SU(2)$ case in [4], some μ -parameters are non-vanishing when $e \neq 0$, even despite the external lines are neglected, therefore, one should not expect the case under consideration makes any sense. However, it turns out in this case still there is a solution of the AGT relation with $\nu \neq 0$:

$$P(a) = a^2 \left(a^4 + 3ea^3 + \frac{13e^2}{4}a^2 + \frac{3e^3}{2}a + \frac{e^4}{4} \right) \tag{83}$$

$$\nu = -\frac{3e^2}{4(-\epsilon_1\epsilon_2)} \tag{84}$$

This polynomial, however, does not correspond to any μ 's.

4.2 Introducing external states

Now we need to switch on non-vanishing $\vec{\alpha}_1, \dots, \vec{\alpha}_4$, and impose the speciality conditions on $\vec{\alpha}_1$ and $\vec{\alpha}_3$, this will modify $B^{(1)}$ at the l.h.s. of (82). Then we use this relation to find the modified polynomial $P(a)$, its roots. Surprisingly or not, they will indeed be just linear functions of α 's and β 's, as required by the AGT conjecture, and they will provide us with the deformation of (44) and (48).

In what follows we omit the normalization factor $-\epsilon_1\epsilon_2$ from most formulas. It can be always restored on dimensional grounds.

The *speciality* condition changes: it is defined by the zeroes of the r.h.s. of (75), and therefore we have six options:

$$\text{either } \beta_1 = \pm Q/2 \quad \text{or} \quad \beta_1 = \pm\alpha_1\sqrt{3} \pm Q \tag{85}$$

for such vectors $\frac{w_1}{\Delta_1} = \sqrt{D_1}$, thus when this ratio appears in our formulas, it does not produce new poles. These six options are of course related by Weyl group transformations. However, since we have two special states in a 4-point conformal block, there is a freedom to choose their relative orientation.

Note that the AGT relation itself is correct for all possible choices in (85). However, in fact, not all of them correspond to the special states (4). Indeed, (4) is more restrictive than just the condition of zero Kac determinant (75). Of 6 states (85) only 3 actually correspond to (4), [5]

$$\text{either } \beta_1 = -Q/2 \quad \text{or} \quad \beta_1 = \pm\alpha_1\sqrt{3} - Q \tag{86}$$

Now all 9 possible combinations of choosing special states $\vec{\alpha}_{1,3}$ should be considered. For instance, instead of (44),

$$\begin{aligned} \mu_1 &= \frac{2}{\sqrt{3}}(\alpha_2 - \alpha_1) = \frac{2}{\sqrt{3}}\alpha_2 - \frac{2\alpha_1}{\sqrt{3}}, \\ \mu_2 &= -\frac{\alpha_2 + 2\alpha_1 + \beta_2\sqrt{3}}{\sqrt{3}} = -\frac{\alpha_2 + \beta_2\sqrt{3}}{\sqrt{3}} - \frac{2\alpha_1}{\sqrt{3}}, \\ \mu_3 &= -\frac{\alpha_2 + 2\alpha_1 - \beta_2\sqrt{3}}{\sqrt{3}} = -\frac{\alpha_2 - \beta_2\sqrt{3}}{\sqrt{3}} - \frac{2\alpha_1}{\sqrt{3}}, \\ \mu_4 &= \frac{2}{\sqrt{3}}(\alpha_4 - \alpha_3) = \frac{2}{\sqrt{3}}\alpha_4 - \frac{2\alpha_3}{\sqrt{3}}, \\ \mu_5 &= -\frac{\alpha_4 + 2\alpha_3 + \beta_4\sqrt{3}}{\sqrt{3}} = -\frac{\alpha_4 + \beta_4\sqrt{3}}{\sqrt{3}} - \frac{2\alpha_3}{\sqrt{3}}, \\ \mu_6 &= -\frac{\alpha_4 + 2\alpha_3 - \beta_4\sqrt{3}}{\sqrt{3}} = -\frac{\alpha_4 - \beta_4\sqrt{3}}{\sqrt{3}} - \frac{2\alpha_3}{\sqrt{3}} \end{aligned} \tag{87}$$

and

$$\nu = -4\alpha_1\alpha_3 \quad (88)$$

one has, e.g., for

$$\underline{\beta_1 = -\epsilon/2, \beta_3 = -\epsilon/2 :}$$

$$\begin{aligned} \mu_1 &= \frac{2\alpha_2 - \alpha_1}{\sqrt{3}} + \frac{1}{2}\epsilon = \frac{2}{\sqrt{3}}\alpha_2 + \frac{1}{\sqrt{3}}\left(-\alpha_1 + \frac{\sqrt{3}}{2}\epsilon\right), \\ \mu_2 &= -\frac{\alpha_2 + \alpha_1 + \beta_2\sqrt{3}}{\sqrt{3}} + \frac{1}{2}\epsilon = -\frac{\alpha_2 + \beta_2\sqrt{3}}{\sqrt{3}} + \frac{1}{\sqrt{3}}\left(-\alpha_1 + \frac{\sqrt{3}}{2}\epsilon\right), \\ \mu_3 &= -\frac{\alpha_2 + \alpha_1 - \beta_2\sqrt{3}}{\sqrt{3}} + \frac{1}{2}\epsilon = -\frac{\alpha_2 - \beta_2\sqrt{3}}{\sqrt{3}} + \frac{1}{\sqrt{3}}\left(-\alpha_1 + \frac{\sqrt{3}}{2}\epsilon\right), \\ \mu_4 &= \frac{2\alpha_4 + \alpha_3}{\sqrt{3}} + \frac{1}{2}\epsilon = \frac{2}{\sqrt{3}}\alpha_4 + \frac{1}{\sqrt{3}}\left(\alpha_3 + \frac{\sqrt{3}}{2}\epsilon\right), \\ \mu_5 &= -\frac{\alpha_4 - \alpha_3 + \beta_4\sqrt{3}}{\sqrt{3}} + \frac{1}{2}\epsilon = -\frac{\alpha_4 + \beta_4\sqrt{3}}{\sqrt{3}} + \frac{1}{\sqrt{3}}\left(\alpha_3 + \frac{\sqrt{3}}{2}\epsilon\right), \\ \mu_6 &= -\frac{\alpha_4 - \alpha_3 - \beta_4\sqrt{3}}{\sqrt{3}} + \frac{1}{2}\epsilon = -\frac{\alpha_4 - \beta_4\sqrt{3}}{\sqrt{3}} + \frac{1}{\sqrt{3}}\left(\alpha_3 + \frac{\sqrt{3}}{2}\epsilon\right) \end{aligned} \quad (89)$$

with

$$\nu = -\left(-\alpha_1 + \frac{\sqrt{3}}{2}\epsilon\right)\left(\alpha_3 + \frac{\sqrt{3}}{2}\epsilon\right) \quad (90)$$

Putting $\alpha_1 = 0$ and $\alpha_3 = 0$ here, one reproduces the solution (84).

Other choices of the special states preserve this structure of answer for μ 's. That is, they are equal to the sum of the same first terms as in (89) plus some quantities $S(\alpha_{1,3})$ which depends on the choice of the special states:

$$\begin{aligned} \mu_1 &= \frac{2}{\sqrt{3}}\alpha_2 + S(\vec{\alpha}_1), & \mu_4 &= -\frac{2}{\sqrt{3}}\alpha_4 + S(\vec{\alpha}_3), \\ \mu_2 &= -\frac{1}{\sqrt{3}}(\alpha_2 + \beta_2\sqrt{3}) + S(\vec{\alpha}_1), & \mu_5 &= \frac{1}{\sqrt{3}}(\alpha_4 + \beta_4\sqrt{3}) + S(\vec{\alpha}_3), \\ \mu_3 &= -\frac{1}{\sqrt{3}}(\alpha_2 - \beta_2\sqrt{3}) + S(\vec{\alpha}_1), & \mu_6 &= \frac{1}{\sqrt{3}}(\alpha_4 - \beta_4\sqrt{3}) + S(\vec{\alpha}_3) \end{aligned} \quad (91)$$

with

$$\begin{aligned} S(\alpha_1, -\epsilon/2) &= -\frac{1}{\sqrt{3}}\alpha_1 + \frac{1}{2}\epsilon, & S(\alpha_1, -\alpha_1\sqrt{3} - \epsilon) &= \frac{2}{\sqrt{3}}\alpha_1 + \epsilon, & S(\alpha_1, \alpha_1\sqrt{3} - \epsilon) &= \frac{2}{\sqrt{3}}\alpha_1 \\ S(\alpha_3, -\epsilon/2) &= \frac{1}{\sqrt{3}}\alpha_3 + \frac{1}{2}\epsilon, & S(\alpha_3, -\alpha_3\sqrt{3} - \epsilon) &= -\frac{2}{\sqrt{3}}\alpha_3, & S(\alpha_3, \alpha_3\sqrt{3} - \epsilon) &= -\frac{2}{\sqrt{3}}\alpha_3 + \epsilon \end{aligned} \quad (92)$$

and

$$\nu = -3S(\alpha_1)S(\alpha_3) \quad (93)$$

The manifest formulas for $S(\alpha_{1,3})$ for all possible choices of the special states are collected in the Conclusion.

5 Calculations at level two

5.1 Level-two Nekrasov functions

Nekrasov functions for $SU(N)$ with $N_f = 2N$ fundamentals – the ones appearing in the AGT representation of the 4-point conformal block with 2 special external states – are labeled by N -plets of ordinary Young diagrams $\vec{Y} = \{Y_1, \dots, Y_N\}$ and are given by [8]:

$$Z_{\vec{Y}} = \frac{\prod_{i=1}^N \prod_{\square \in Y_i} \prod_{f=1}^{N_f} \left(\phi(a_i, Y_i, \square) + \mu_f \right)}{\prod_{i,j=1}^N \prod_{\square \in Y_i} E(a_i - a_j, Y_i, Y_j, \square) \left(\epsilon - E(a_i - a_j, Y_i, Y_j, \square) \right)} \quad (94)$$

where for a box with coordinates (m, n) in the Young diagram

$$\begin{aligned} \phi(a, Y, \square_{m,n}) &= a + \epsilon_1(m-1) + \epsilon_2(n-1), \\ E(a, Y_1, Y_2, \square_{m,n}) &= a + \epsilon_1 \left(k_n^T(Y_1) - m + 1 \right) - \epsilon_2 \left(k_m(Y_2) - n \right) \end{aligned} \quad (95)$$

$k_n^T(Y)$ and $k_m(Y)$ are the length of the n -th row and the height of the m -th column in the diagram Y .

If $Y_2 = Y_1$, then at $\epsilon = 0$ the combination $h(\square_{m,n}) = E(0, Y, Y) = (k_n^T(Y) - m + 1) + ((k_m(Y) - n))$ is the hook length for the box (m, n) , which enters, for example, the celebrated hook formula for

$$d_Y = \prod_{\square \in Y} h^{-1}(\square) \quad (96)$$

for the character $\chi_Y(t_k = \delta_{k,1}) = d_Y$, which defines up to various simple factors dimensions of representations of symmetric and linear groups. Thus, for $\epsilon = 0$ the Nekrasov formulas are more or less natural objects in the theory of character expansions [28, 29] and Hurwitz-Kontsevich partition functions [30], while for $\epsilon \neq 0$ they belong to representation theory of more sophisticated (quantum affine?) algebras. The AGT relation associates the deformation to $\epsilon \neq 0$ with the deformation to $c \neq r = N - 1$ in the theory of r free fields (conformal Toda?). The square d_Y^2 is also known as the Plancherel measure for integer partitions, which appears in the theory of KP and Kontsevich-Hurwitz τ -functions, beginning from the famous sum rule

$$\sum_Y \chi_Y(t) \chi_Y(t') = \exp \left(\sum_k k t_k t'_k \right) \quad (97)$$

and its particular case

$$\sum_Y d_Y^2 x^{|Y|} = e^x \quad (98)$$

The Nekrasov formulas with $\epsilon \neq 0$ provide a deformation of this measure.

At level one, only one kind of N -plets $\vec{Y}$ contributes, containing just one diagram of the unit size:

$$Z_{[1]} = -\frac{1}{\epsilon_1 \epsilon_2} \sum_{i=1}^N \frac{P(a_i)}{\prod_{k \neq i}^N a_{ik}(a_{ik} + \epsilon)} \equiv -\frac{1}{\epsilon_1 \epsilon_2} \sum_{i=1}^N R_i(a_i) \quad (99)$$

At level two, only three kinds of N -plets $\vec{Y}$ contribute: containing just one diagram of size two, either [2] or [11], or containing a pair of the single-box [1] diagrams. The corresponding Nekrasov functions are

$$\begin{aligned} Z_{[2]} &= \frac{1}{2! \epsilon_1 \epsilon_2^2 (\epsilon_1 - \epsilon_2)} \sum_{i=1}^N R_i(a_i) R_i(a_i + \epsilon_2), \\ Z_{[11]} &= -\frac{1}{2! \epsilon_1^2 \epsilon_2 (\epsilon_1 - \epsilon_2)} \sum_{i=1}^N R_i(a_i) R_i(a_i + \epsilon_1), \\ Z_{[1][1]} &= \frac{1}{\epsilon_1^2 \epsilon_2^2} \sum_{i < j}^N R_i(a_i) R_j(a_j) \frac{a_{ij}^2 (a_{ij}^2 - \epsilon^2)}{(a_{ij}^2 - \epsilon_1^2)(a_{ij}^2 - \epsilon_2^2)} \end{aligned} \quad (100)$$

See also eq.(62) of [4] for a particular case of $N = 2$.

5.2 AGT relations at level two

The AGT relation at level two states that

$$B^{(2)} = Z_{[2]} + Z_{[11]} + Z_{[1][1]} + \nu Z_{[1]} + \frac{\nu(\nu + 1)}{2} \quad (101)$$

where $Z_{[1]} = Z_1$. The denominator at the l.h.s. comes from inverse of the Shapovalov form, i.e. equals to the Kac determinant of matrix (65), while at the r.h.s. it contains the product of a_{ij} , shifted by linear combinations of ϵ_1 and ϵ_2 . In other words, the AGT relation implies, to begin with, the matching of the denominators, which requires that

$$\text{Kac determinant} = \det \left(\text{Shapovalov matrix} \right) \sim \prod_{i \neq j} \prod_{p, q > 0} (a_{ij} + p\epsilon_1 + q\epsilon_2) \quad (102)$$

The fact that the Kac determinants factorize in such an elegant way, if expressed through the $\vec{\alpha}$ -variables instead of the eigenvalues Δ and w of the W -operators, is one of the central results of representation theory of the Virasoro and W -algebras. In s.5.3 we illustrate this remarkable general result in the particular example of level two for the W_3 algebra.

Note, however, that the sets $\{(p, q)\}$ are actually not obviously the same at the l.h.s. and the r.h.s. of (102). In the Kac determinant they are constrained by the condition $0 < pq \leq \text{level}$, while in Nekrasov's formulas at the same level one can expect that either p , or q can be equal to zero. These extra zeroes of the Nekrasov determinant are actually canceled between different terms in the sum over Young diagrams, however, particular Nekrasov functions have *more* singularities than there are zeroes of the Kac determinant.

An additional claim as compared to the Kac determinant factorization property in the AGT conjecture is that the numerators also match for an appropriate *linear* relation between μ_f and $\vec{a}_{1,2,3,4}$ parameters (with $\vec{a}_{1,3}$ subjected to the *speciality* constraints). We do not present details of this check here: at the present level of understanding it is more a computer exercise than a conceptual calculation. Some necessary comments are given in s.5.4.

5.3 Factorization of Kac determinant

For $c = 2$ the determinants of the level-one and level-two Shapovalov matrices (11) and (65) are equal to

$$3^2 \beta^2 (\beta^2 - 3\alpha^2)^2 \sim \left(a_{12} a_{23} a_{31} \right)^2 \quad (103)$$

and

$$\begin{aligned} & 3^8 \beta^4 (2\beta - 1)^2 (2\beta + 1)^2 (\beta^2 - 3\alpha^2)^4 \left((\beta - 1)^2 - 3\alpha^2 \right)^2 \left((\beta + 1)^2 - 3\alpha^2 \right)^2 \sim \\ & \sim \left((a_{12} - 1) a_{12}^2 (a_{12} + 1) (a_{23} - 1) a_{23}^2 (a_{23} + 1) (a_{31} + 1) a_{31}^2 (a_{31} - 1) \right)^2 \end{aligned} \quad (104)$$

respectively. This is in perfect agreement with (102), provided $\epsilon_1 = -\epsilon_2 = 1$ and identification (23) is made:

$$\begin{aligned} & a_{12} = -2\beta, \quad a_{23} = \beta + \alpha\sqrt{3}, \quad a_{31} = \beta - \alpha\sqrt{3}, \quad \text{i.e.} \\ & a_1 = \frac{1}{\sqrt{3}}(\alpha - \beta\sqrt{3}), \quad a_2 = \frac{1}{\sqrt{3}}(\alpha + \beta\sqrt{3}), \quad a_3 = -\frac{2\alpha}{\sqrt{3}} \end{aligned} \quad (105)$$

(the relative normalization of a and α in (105) is defined by the full AGT relation and is not obvious at the level of Kac determinants).

For generic c the degeneration that gives rise to full squares in the Nekrasov denominators is resolved, and the same happens to the Kac determinant. At level one, one obtains instead of (103):

$$\begin{aligned} K_1 &= \frac{3^2}{4 - 15Q^2} (4\beta^2 - Q^2) \left((\beta - Q)^2 - 3\alpha^2 \right) \left((\beta + Q)^2 - 3\alpha^2 \right) \sim \\ & \sim (2\beta - Q)(2\beta + Q)(\beta + \alpha\sqrt{3} - Q)(\beta + \alpha\sqrt{3} + Q)(\beta - \alpha\sqrt{3} - Q)(\beta - \alpha\sqrt{3} + Q) \sim \\ & \sim \prod_{i \neq j} (a_{ij} + \epsilon) = (a_{12}^2 - \epsilon^2)(a_{23}^2 - \epsilon^2)(a_{31}^2 - \epsilon^2) \end{aligned} \quad (106)$$

Therefore, identification (105) remains intact, one should only add that $Q = \epsilon$.

At level two, the l.h.s. of (104) is deformed into

$$\begin{aligned} K_2 &= \frac{2^4 \cdot 3^8}{(4 - 15Q^2)^4} \left\{ (4\beta^2 - Q^2) \left((\beta - Q)^2 - 3\alpha^2 \right) \left((\beta + Q)^2 - 3\alpha^2 \right) \right\}^2 \cdot \\ & \quad (2Q^2 - 1 - 6\beta Q + 4\beta^2)(2Q^2 - 1 + 6\beta Q + 4\beta^2) \cdot \\ & \quad (1 + 12\beta Q^3 + 6\beta^3 Q - 6\beta Q - 18\alpha^2 \beta Q - 15\alpha^2 Q^2 - 6\alpha^2 - 2\beta^2 - 4Q^2 - 6\alpha^2 \beta^2 + 13\beta^2 Q^2 + \beta^4 + 4Q^4 + 9\alpha^4) \cdot \\ & \quad (1 - 12\beta Q^3 - 6\beta^3 Q + 6\beta Q + 18\alpha^2 \beta Q - 15\alpha^2 Q^2 - 6\alpha^2 - 2\beta^2 - 4Q^2 - 6\alpha^2 \beta^2 + 13\beta^2 Q^2 + \beta^4 + 4Q^4 + 9\alpha^4) \end{aligned} \quad (107)$$

so that the level one Kac determinant factors out from that of level two for arbitrary Δ and w , and the rest of the formula (including the second power of (106)) factorizes nicely in the α -parametrization. The dictionary for decoding this quantity is simple: for $\epsilon_1 + \epsilon_2 = \epsilon = Q$ and $\epsilon_1 \epsilon_2 = -1$ one has

$$\begin{aligned} & (2\beta + \epsilon_1)(2\beta + \epsilon_2) = 4\beta^2 + 2Q\beta - 1, \\ & (2\beta + 2\epsilon_1 + \epsilon_2)(2\beta + \epsilon_1 + 2\epsilon_2) = 4\beta^2 + 6Q\beta + 2Q^2 - 1 \end{aligned} \quad (108)$$

and

$$\begin{aligned} & (\beta + \alpha\sqrt{3} + \epsilon_1)(\beta + \alpha\sqrt{3} + \epsilon_2)(\beta - \alpha\sqrt{3} + \epsilon_1)(\beta - \alpha\sqrt{3} + \epsilon_2) = \\ & 1 - 6\alpha^2 \beta Q + 2Q\beta^3 + 2Q\beta - 6\alpha^2 \beta^2 - 3\alpha^2 Q^2 + 6\alpha^2 + 2\beta^2 + \beta^2 Q^2 + 9\alpha^4 \beta^4, \\ & (\beta + \alpha\sqrt{3} + 2\epsilon_1 + \epsilon_2)(\beta + \alpha\sqrt{3} + \epsilon_1 + 2\epsilon_2)(\beta - \alpha\sqrt{3} + 2\epsilon_1 + \epsilon_2)(\beta - \alpha\sqrt{3} + \epsilon_1 + 2\epsilon_2) = \\ & = (1 + 12\beta Q^3 + 6\beta^3 Q - 6\beta Q - 18\alpha^2 \beta Q - 15\alpha^2 Q^2 - 6\alpha^2 - 2\beta^2 - 4Q^2 - 6\alpha^2 \beta^2 + 13\beta^2 Q^2 + \beta^4 + 4Q^4 + 9\alpha^4) \end{aligned} \quad (109)$$

Thus, one can see that

$$K_2 \sim (a_{12}^2 - \epsilon^2)^2 (a_{23}^2 - \epsilon^2)^2 (a_{31}^2 - \epsilon^2)^2 \left(a_{12}^2 - (\epsilon_1 + 2\epsilon_2)^2 \right) \left(a_{23}^2 - (\epsilon_1 + 2\epsilon_2)^2 \right) \left(a_{31}^2 - (\epsilon_1 + 2\epsilon_2)^2 \right) \cdot \left(a_{12}^2 - (2\epsilon_1 + \epsilon_2)^2 \right) \left(a_{23}^2 - (2\epsilon_1 + \epsilon_2)^2 \right) \left(a_{31}^2 - (2\epsilon_1 + \epsilon_2)^2 \right) \quad (110)$$

The factors $(a_{ij}^2 - \epsilon_1^2)$ and $(a_{ij}^2 - \epsilon_2^2)$, which one could expect to arise looking at the denominators in eq.(100), do not actually appear in the Kac determinant. Instead, K_1 comes squared. This means that these factors should cancel out after summing different terms in the Nekrasov formula with the appropriately chosen polynomial $P(a)$ in the numerator.

5.4 Conformal block at level two

The conformal block $B^{(2)}$ is obtained from the general expression (10) by substitution of the inverted 5×5 Shapovalov matrix (65) and five pairs triple vertices from [5]. To define these vertices unambiguously in unspecified conformal model, one should impose the *speciality* conditions and make use of (62), what is also done in [5].

Note that there are five generalized Young diagrams $\mathcal{Y}$ of size $|\mathcal{Y}| = 2$, two "pure Virasoro", two "pure W " and one "mixed". Starting from level two, there is no way to separate W and Virasoro diagrams. Collecting all the seven pairs of vertices for the two first levels in one place, one gets

$\begin{aligned} \bar{\Gamma}(L_{-1}) &= \Delta + \Delta_1 - \Delta_2, \\ \bar{\Gamma}(W_{-1}) &= w + 2w_1 - w_2 + \frac{3w_1}{2\Delta_1}(\Delta - \Delta_1 - \Delta_2) \end{aligned}$	(111)
$\begin{aligned} \bar{\Gamma}(L_{-2}) &= \Delta + 2\Delta_1 - \Delta_2, \\ \bar{\Gamma}(L_{-1}^2) &= (\Delta + \Delta_1 - \Delta_2)(\Delta + \Delta_1 - \Delta_2 + 1), \\ \bar{\Gamma}(L_{-1}W_{-1}) &= (\Delta + \Delta_1 - \Delta_2 + 1) \left(w + 2w_1 - w_2 + \frac{3w_1}{\Delta_1}(\Delta - \Delta_1 - \Delta_2) \right), \\ \bar{\Gamma}(W_{-2}) &= w + 5w_2 - w_2 + \frac{3w_1}{\Delta_1}(\Delta - \Delta_1 - \Delta_2), \\ \bar{\Gamma}(W_{-1}^2) &= \left(w + 2w_1 - w_2 + \frac{3w_1}{2\Delta_1}(\Delta - \Delta_1 - \Delta_2) \right) \left(w + 2w_1 - w_2 + \frac{3w_1}{2\Delta_1}(\Delta - \Delta_1 - \Delta_2 + 1) \right) + \\ &\quad + \frac{9D}{2}(\Delta + \Delta_1 - \Delta_2) \end{aligned}$	
$\begin{aligned} \Gamma(L_{-1}) &= \Delta + \Delta_3 - \Delta_4, \\ \Gamma(W_{-1}) &= w + w_3 + w_4 - \frac{3w_3}{2\Delta_3}(\Delta + \Delta_3 - \Delta_4) \end{aligned}$	
$\begin{aligned} \Gamma(L_{-2}) &= \Delta + 2\Delta_3 - \Delta_4, \\ \Gamma(L_{-1}^2) &= (\Delta + \Delta_3 - \Delta_4)(\Delta + \Delta_3 - \Delta_4 + 1), \\ \Gamma(L_{-1}W_{-1}) &= (\Delta + \Delta_3 - \Delta_4 + 1) \left(w + w_3 + w_4 - \frac{3w_3}{\Delta_3}(\Delta + \Delta_3 - \Delta_4) \right), \\ \Gamma(W_{-2}) &= w + w_3 + w_4 - \frac{3w_3}{\Delta_3}(\Delta + \Delta_3 - \Delta_4), \\ \Gamma(W_{-1}^2) &= \left(w + w_3 + w_4 - \frac{3w_3}{2\Delta_3}(\Delta + \Delta_3 - \Delta_4) \right) \left(w + w_3 + w_4 - \frac{3w_3}{2\Delta_3}(\Delta + \Delta_3 - \Delta_4 + 1) \right) + \\ &\quad + \frac{9D}{2}(\Delta + \Delta_3 - \Delta_4) \end{aligned}$	

Here we assumed that $\vec{\alpha}_1$ and $\vec{\alpha}_3$ are the special state, i.e. that

$$W_{-1}V_{\alpha_1}(1) = \frac{3w_1}{\Delta_1}L_{-1}V_{\alpha_1}(1) \stackrel{(59)}{=} -\frac{3w_1}{2\Delta_1}(\Delta + \Delta_1 - \Delta_2)V_{\alpha_1}(1) \quad (112)$$

and similarly for α_3 : this is why these specific combinations appear in the formulas. We also omit the common structure constant factors $C_{\alpha_1\alpha_2}^\alpha = \langle V_\alpha(0)|V_{\alpha_1}(1)V_{\alpha_2}(\infty)\rangle$ and $C_{\alpha\alpha_3\alpha_4} = \langle V_\alpha(0)V_{\alpha_1}(1)V_{\alpha_2}(\infty)\rangle$ at the r.h.s. Note the delicate and seemingly irregular sign differences between formulas for $\bar{\Gamma}$ and Γ vertices, all these details being essential, since the AGT relations are very sensitive to details.

For making up a conformal block, one also needs to invert the Shapovalov matrices (64) and (65). For (64) it is simple:

$$\frac{1}{K_1} \begin{vmatrix} 9D\Delta/2 & -3w \\ -3w & 2\Delta \end{vmatrix} \quad (113)$$

but this paper is too short to explicitly write down the inverse of (65), neither in Δ, w , nor in α, β variables. Still, multiplying this inverse by the two 5-vector Γ and $\bar{\Gamma}$ from (111) accordingly to (10), one obtains the conformal block $B^{(2)}$ which, indeed, coincides with the combination of $Z^{(2)}$ and $Z^{(1)}$ at the r.h.s. of (101), provided the six μ 's and ν are given by relations (91)-(93) for various choices of *special* states $\vec{\alpha}_1$ and $\vec{\alpha}_3$, which we already found explicitly in the analysis of the first level. The calculation is tedious, it necessarily includes the check of expressions (111) for the 3-point functions. As usual, it is worth starting from the simplest case of $c = 2$, $\vec{\alpha}_1 = \dots = \vec{\alpha}_4 = 0$ (for $c = 2$ this does not contradict *speciality* conditions). In this case, (111) is simplified to

$\Gamma(L_{-1}) = \Delta$	$= \Delta,$
$\Gamma(W_{-1}) = w$	$= w$
$\Gamma(L_{-2}) = \Delta$	$= \Delta,$
$\Gamma(L_{-1}^2) = \Delta(\Delta + 1)$	$= \Delta(\Delta + 1),$
$\Gamma(L_{-1}W_{-1}) = (\Delta + 1)w$	$= (\Delta + 1)w,$
$\Gamma(W_{-2}) = x_1w$	$= w,$
$\Gamma(W_{-1}^2) = x_2\Delta^3 + x_3w^2 + x_4\Delta^2 + x_5\Delta$	$= w^2 + \frac{9}{2}\Delta^2$
$\Gamma(L_{-1}) = \Delta$	$= \Delta,$
$\Gamma(W_{-1}) = w$	$= w$
$\Gamma(L_{-2}) = \Delta$	$= \Delta,$
$\Gamma(L_{-1}^2) = \Delta(\Delta + 1)$	$= \Delta(\Delta + 1),$
$\Gamma(L_{-1}W_{-1}) = (\Delta + 1)w$	$= (\Delta + 1)w,$
$\Gamma(W_{-2}) = y_1w$	$= w,$
$\Gamma(W_{-1}^2) = y_2\Delta^3 + y_3w^2 + y_4\Delta^2 + y_5\Delta$	$= w^2 + \frac{9}{2}\Delta^2$

(114)

and the Nekrasov formula can be expressed through Δ and w as follows:

$$Z^{(2)} = \frac{2}{K_2} \left\{ 81\Delta(\Delta^3 - w^2) \left(\Delta^3 - \frac{8}{9}w^2 \right) \right\}^2 +$$

$$\begin{aligned}
& + \frac{81}{4K_2}(\Delta^2 - w^2) \left\{ \Delta \left(-1863\Delta^9 + 2160w^2\Delta^6 + 608w^4\Delta^3 - 896w^6 \right) + 6 \left(81\Delta^9 + 480w^2\Delta^6 - 448w^4\Delta^3 \right) + \right. \\
& + \Delta^2 \left(3807\Delta^6 - 6696w^2\Delta^3 + 2704w^4 \right) + 4\Delta \left(-1377\Delta^6 + 1422w^2\Delta^3 - 152w^4 \right) + 3 \left(1053\Delta^6 - 960w^2\Delta^3 + 16w^4 \right) + \\
& \left. - 81\Delta(10\Delta - 1) \left(\Delta^3 - \frac{8}{9}w^2 \right) \right\} = \tag{115}
\end{aligned}$$

$$\begin{aligned}
& = \frac{81\Delta^4(\Delta - 1)^4 \left(8\Delta^3 + 9\Delta^2 - 6\Delta + 1 \right) + G_2w^2 + G_3w^4 + G_4w^6}{4 \cdot 81 \cdot (\Delta^3 - w^2) \left(4\Delta(\Delta - 1)^2 - 4w^2 \right)^2}, \\
& G_2 = -72\Delta(\Delta - 1)^2(25\Delta^5 + 20\Delta^4 - 25\Delta^3 + 23\Delta^2 - 8\Delta + 1), \\
& G_4 = 1664\Delta^5 + 608\Delta^4 - 2688\Delta^3 + 2704\Delta^2 - 608\Delta + 48, \\
& G_6 = -128\Delta(4\Delta + 7), \tag{116}
\end{aligned}$$

where we grouped terms with the same power of $-\epsilon_1\epsilon_2$ (which is suppressed for sake of space, but can be restored in the formulas) or, alternatively, with the same power of w .

Eq.(101) with $P(a) = a^6$, $\nu = 0$ and AGT relation (23) between $\vec{a}$ and $\vec{\alpha}$ can be *used* to find the ten free parameters $x_1, \dots, y_5$. It is a strongly over-defined system of equations, still it has a solution: $x_1 = 1$, $x_2 = 0$, $x_3 = 1$, $x_4 = 9/2$, $x_5 = 0$, $y_1 = 1$, $y_2 = 0$, $y_3 = 1$, $y_4 = 9/2$, $y_5 = 0$, as shown in the right column of the table (114). Then, validation of the AGT relation at this level comes from comparison of *such* x 's and y 's with their values in the CFT table (111). After that, one switches on the external states $\vec{\alpha}_1, \dots, \vec{\alpha}_4$ and finally deforms to $c \neq 2$. **This completes the explicit check of the W_3 AGT relation at level two.**

6 On complete proof of AGT relation in a *very special* case

In [6], among other things, we provided a proof of the AGT relation in a very restricted setting. The idea (also mentioned in [2]) is to make use of exact knowledge of the 4-point conformal block in the $sl(N)$ Toda theory [17] for the *very special* kinematics: when one of external lines is *special* and another one is further restricted to belong to the *fully degenerate* W -Verma module. In $W^{(N)}$ case *special* means that $N - 2$ conditions of the type (4) are imposed on the $(N - 1)$ -component momentum, let it be $\vec{\alpha}_3$, while *fully degenerate* means that the last remaining component of the *special* $\vec{\alpha}_1$ is further fixed to a certain value. Selection rules of the Toda theory expresses the internal-state momentum $\vec{\alpha}$ through $\vec{\alpha}_1$ and $\vec{\alpha}_2$, much similarly to the free field model rule $\vec{\alpha} = \vec{\alpha}_1 + \vec{\alpha}_2$. The only difference is that, in the Toda case, there are not single, but N possible values of $\vec{\alpha}$ (i.e. N different non-zero conformal blocks). Thus, this Fateev-Litvinov conformal block depends on $2(N - 1) + 1 + 1$ free parameters ($\vec{\alpha}_2, \alpha_3, \vec{\alpha}_4, c$) and on this high-codimension subspace in the total moduli space of the 4-point conformal blocks it is represented by *generic* hypergeometric series

$$\begin{aligned}
{}_N F_{N-1}(A_1, \dots, A_N; B_1, \dots, B_{N-1}, x) &= 1 + x \cdot \frac{A_1 \dots A_N}{B_1 \dots B_{N-1}} + \frac{x^2}{2} \cdot \frac{A_1(A_1 + 1) \dots A_N(A_N + 1)}{B_1(B_1 + 1) \dots B_N B_{N+1}} + \\
&+ \sum_{n=3}^{\infty} \frac{x^n}{n!} \cdot \prod_{k=0}^{n-1} \frac{(A_1 + k) \dots (A_N + k)}{(B_1 + k) \dots (B_{N-1} + k)} \tag{117}
\end{aligned}$$

where $2N - 1$ parameters A_i, B_i are linear combinations of $\vec{\alpha}_2, \alpha_3, \vec{\alpha}_4$ with coefficients, depending on the central charge (actually, on the screening-charge parameters ϵ_1 and ϵ_2). This remarkable result demonstrates how generic hypergeometric series are embedded, as a linear subspace if the α -parametrization of this moduli space is used!, into the space of conformal blocks, thus, generalizing the old description [27] of their embedding into the space of Dotsenko-Fateev integrals, or of hypergeometric integrals/correlators in the terminology of [26] (we remind that arbitrary ${}_N F_{N-m}$ are obtained in certain limits from ${}_N F_{N-1}$ and lie on the boundary of the moduli space, see, for example, [27]). As explained in [6], in these terms the AGT conjecture states that **the Nekrasov functions provide the generalization of the $N = 2$ hypergeometric series, exactly extending it to entire space of Virasoro conformal blocks**. For $N > 2$ they are probably sufficient only to extend hypergeometric series to the moduli space of conformal blocks, restricted to *special* subspace (i.e. when only two out of m external momenta and all the $m - 3$ internal momenta in the m -point conformal block are arbitrary, while the other $m - 2$ external momenta belong to 1-dimensional *special* subspaces). If this is true, there remains a question, **what provides the further extension beyond Nekrasov functions(?)**, which still remains to be answered.

However, if one reverse the question: what are hypergeometric series from the point of view of Nekrasov functions, one immediately arrives to complete proof of the AGT conjecture for this very restricted (still huge!) class of conformal blocks. The point is that the answer is very simple: the hypergeometric series arise when only very specific N -plets of Young diagrams are non-vanishing: $\vec{Y} = \{\emptyset, \dots, [1^n], \dots, \emptyset\}$ or $\vec{Y} = \{\emptyset, \dots, [n], \dots, \emptyset\}$, i.e. only one diagram in the N -plet is not empty, and it is either a row $[1^n]$ or column $[n]$. We call such Nekrasov functions *chiral* and *anti-chiral* respectively, explicitly they are

$$Z(\emptyset, \dots, [1^n], \dots, \emptyset) = \frac{1}{\epsilon_2 \epsilon_1^n n!} \frac{P(a_i) P(a_i + \epsilon_1) \dots P(a_i + (n-1)\epsilon_1)}{(-\epsilon_2)(-\epsilon_2 + \epsilon_1) \dots (-\epsilon_2 + (n-1)\epsilon_1) Q_i(a_i) Q_i(a_i + \epsilon_1) \dots Q_i(a_i + (n-1)\epsilon_1)} \quad (118)$$

and

$$Z(\emptyset, \dots, [n], \dots, \emptyset) = Z(\emptyset, \dots, [1^n], \dots, \emptyset) (\epsilon_1 \leftrightarrow \epsilon_2) \quad (119)$$

where i is the position of the non-empty diagram in the N -plet, $Q_i(x) = \prod_{j \neq i} (x - a_j)(x - a_j + \epsilon)$ and these functions immediately reproduce the hypergeometric series = the Fateev-Litvinov conformal block (117):

$$P_i(a_i) = \epsilon_1^N A_1 \dots A_N, \quad Q_i(a_i) = \frac{\epsilon_1^N B_1 \dots B_{N-1}}{-\epsilon_1 \epsilon_2} \quad (120)$$

for the chiral function. However, there is a question: what guarantees that only the chiral functions are non-vanishing? As explained in [6], this is a condition on polynomial P : for given i ,

$$P_i(a_j) = 0 \quad \text{for } j \neq i \quad \text{and} \quad P_i(a_i + \epsilon_2) = 0 \quad (121)$$

These are, in fact, N conditions on the coefficients (or on the parameters μ_f) of the polynomial $P(a) = \prod_{f=1}^{N_f=2N} (a + \mu_f)$ of degree $2N$, and the remaining N parameters can be used to match the arbitrary set of $A_1, \dots, A_N$ in (120). After that the $N-1$ parameters a_j ($\sum_{j=1}^N a_j = 0$) can be adjusted to match $B_1, \dots, B_{N-1}$. Note that all these relations are *linear*, in accordance with the general AGT claim. The parameters ϵ_1 and ϵ_2 are associated with the central charge, and the common scale drops out from the Nekrasov functions in the case of $N_f = 2N$, as usual.

We now use the detailed description of the $N = 3$ case in this paper to show explicitly how these matchings work in this particular case, and illustrate what (121) has to do with the maximal degeneracy condition imposed in [17]. The simplest is to pick up formulas from the Conclusion, where they are collected in a list. For the sake of definiteness, we choose $i = 3$ in (121) (the choice of i corresponds to the choice of non-zero conformal block, there are exactly N of them). Then, it is obvious from (130) and (131) that there is a natural choice of solution to $P(a_1) = P(a_2) = 0$:

$$\begin{aligned} 0 = (a_1 + \mu_3) &= \frac{1}{\sqrt{3}}(\alpha - \alpha_2) - (\beta - \beta_2) + S(\vec{\alpha}_1), \\ 0 = (a_2 + \mu_2) &= \frac{1}{\sqrt{3}}(\alpha - \alpha_2) + (\beta - \beta_2) + S(\vec{\alpha}_1) \end{aligned} \quad (122)$$

i.e.

$$\alpha = \alpha_2 - \sqrt{3}S(\vec{\alpha}_1), \quad \beta = \beta_2 \quad (123)$$

which restricts the intermediate momentum $\vec{\alpha}$ to be a special combination of the two external momenta $\vec{\alpha}_1$ and $\vec{\alpha}_2$. The third condition $P(a_3 + \epsilon_1) = 0$ is then naturally imposed as a constraint

$$0 = ((a_3 + \epsilon_1) + \mu_1) = -\frac{2}{\sqrt{3}}(\alpha - \alpha_2) + \epsilon_1 + S(\vec{\alpha}_1) \stackrel{(123)}{=} 3S(\vec{\alpha}_1) + \epsilon_1 \quad (124)$$

which imposes an additional restriction on the already-*special* momentum $\vec{\alpha}_1$, i.e. fixes it completely. With these choices, (123) and (124) it is guaranteed that the Nekrasov partition function is an $N = 3$ hypergeometric series (117) with

$$\begin{aligned} P(a_3) &= \prod_{f=1}^6 (a_3 + \mu_f) \stackrel{(123) \& (124)}{=} (-\epsilon_1) \left((\epsilon + \sqrt{3}\alpha_2)^2 - \beta_2^2 \right) \prod_{f=4}^6 (a_3 + \mu_f) = \\ &= (-\epsilon_1) \left((\epsilon + \sqrt{3}\alpha_2)^2 - \beta_2^2 \right) A_1 A_2 A_3 \end{aligned} \quad (125)$$

and

$$a_{31}a_{32}(a_{31} + \epsilon)(a_{32} + \epsilon) \stackrel{(123)\&(124)}{=} \left((\epsilon + \sqrt{3}\alpha_2)^2 - \beta_2^2 \right) (a_{31} + \epsilon)(a_{32} + \epsilon) = \left((\epsilon + \sqrt{3}\alpha_2)^2 - \beta_2^2 \right) B_1 B_2 \quad (126)$$

i.e.

$$A_s = a_3 + \mu_{s+3} = \left\{ -\frac{2}{\sqrt{3}}\alpha_2 - \frac{2}{3}\epsilon + \mu_4, -\frac{2}{\sqrt{3}}\alpha_2 - \frac{2}{3}\epsilon + \mu_5, -\frac{2}{\sqrt{3}}\alpha_2 - \frac{2}{3}\epsilon + \mu_6 \right\}, \quad s = 1, 2, 3, \\ B_s = a_{3s} + \epsilon = \left\{ \sqrt{3}\alpha_2 - \beta_2, \sqrt{3}\alpha_2 + \beta_2 \right\}, \quad s = 1, 2 \quad (127)$$

with $\mu_{4,5,6}$ from (131).

It remains to say that (124) is exactly the condition for $\vec{\alpha}_1$ to describe the maximally degenerate state, and (123) is exactly the relevant selection rule in the $sl(3)$ Toda model [17]. **This completes the proof of AGT conjecture for the generic $N = 3$ hypergeometric conformal blocks. The proof remains the same for arbitrary N ,** provided we accept the straightforward generalization (138) of (130) and (131). A similarly transparent proof for non-hypergeometric blocks, however, remains to be found.

7 Summary and conclusion

Consideration in this paper leaves **little room for doubts that the AGT relations are true for the Verma modules of generic groups**, *provided appropriate speciality* (null-vector) constraints are imposed on the modules. Exact linear expressions for the Nekrasov parameters for the 4-point conformal block are³:

$$SU(2) : \quad a_1 = -a_2 = \alpha, \quad \mu_{1,2} = \pm\alpha_2 + S(\alpha_1), \quad \mu_{3,4} = \pm\alpha_4 + S(\alpha_3), \quad \nu = \frac{2S(\alpha_1)S(\alpha_2)}{\epsilon_1\epsilon_2} \quad (128)$$

for the $SU(2)$ case. There are two different choices for each of the two shifts $S(\alpha_1)$ and $S(\alpha_3)$ (and thus a total of 4 choices for the pair),

$$S(\pm\alpha) = \frac{\epsilon}{2} \pm \alpha \quad (129)$$

Similarly, for $SU(3)$

$$\boxed{a_1 = \frac{1}{\sqrt{3}}(\alpha - \beta\sqrt{3}), \quad a_2 = \frac{1}{\sqrt{3}}(\alpha + \beta\sqrt{3}), \quad a_3 = -\frac{2\alpha}{\sqrt{3}}} \quad (130)$$

$$\boxed{\begin{aligned} \mu_1 &= \frac{2}{\sqrt{3}}\alpha_2 + S(\vec{\alpha}_1), & \mu_4 &= -\frac{2}{\sqrt{3}}\alpha_4 + S(\vec{\alpha}_3), \\ \mu_2 &= -\frac{1}{\sqrt{3}}(\alpha_2 + \beta_2\sqrt{3}) + S(\vec{\alpha}_1), & \mu_5 &= \frac{1}{\sqrt{3}}(\alpha_4 + \beta_4\sqrt{3}) + S(\vec{\alpha}_3), \\ \mu_3 &= -\frac{1}{\sqrt{3}}(\alpha_2 - \beta_2\sqrt{3}) + S(\vec{\alpha}_1), & \mu_6 &= \frac{1}{\sqrt{3}}(\alpha_4 - \beta_4\sqrt{3}) + S(\vec{\alpha}_3) \end{aligned}} \quad (131)$$

$$\boxed{\nu = \frac{3S(\vec{\alpha}_1)S(\vec{\alpha}_3)}{\epsilon_1\epsilon_2}} \quad (132)$$

There are three different choices for each of the two shifts $S(\vec{\alpha}_1)$ and $S(\vec{\alpha}_3)$ (and thus a total of 9 choices for the pair), associated with the three possibilities to choose a *special* values⁴ $\vec{\alpha}_1 = (\alpha_1, \beta_1)$ and $\vec{\alpha}_3 = (\alpha_3, \beta_3)$:

$$\boxed{S(\alpha_1, -\epsilon/2) = -\frac{1}{\sqrt{3}}\alpha_1 + \frac{1}{2}\epsilon, \quad S(\alpha_1, -\alpha_1\sqrt{3} - \epsilon) = \frac{2}{\sqrt{3}}\alpha_1 + \epsilon, \quad S(\alpha_1, \alpha_1\sqrt{3} - \epsilon) = \frac{2}{\sqrt{3}}\alpha_1} \quad (133)$$

$$\boxed{S(\alpha_3, -\epsilon/2) = \frac{1}{\sqrt{3}}\alpha_3 + \frac{1}{2}\epsilon, \quad S(\alpha_3, -\alpha_3\sqrt{3} - \epsilon) = -\frac{2}{\sqrt{3}}\alpha_3, \quad S(\alpha_3, \alpha_3\sqrt{3} - \epsilon) = -\frac{2}{\sqrt{3}}\alpha_3 + \epsilon} \quad (134)$$

³Note that as compared with [4], our definition of α is shifted: $\alpha \rightarrow \alpha + \frac{\epsilon}{2}$.

⁴We define these *special values* simply as zeroes of the Kac determinant $\bar{K}_1$, i.e. from the condition $D\Delta^2 = w^2$. Actual *special states*, satisfying (4) do not obligatory exist for each of these values (see, for example, the free-model example in [5], where only three out of six possible *special values* are actually associated with *special states*). It turns out that for this does not matter for AGT relation: it holds just for all *special values*.

If some other pair of external states is chosen to be *special*, formulas change according to the rule of projective transformation of the conformal block. The resulting ν can be easily evaluated:

$$1, 3 \leftrightarrow 2, 4 \quad \nu \rightarrow \frac{3S(\vec{\alpha}_2)S(\vec{\alpha}_4)}{\epsilon_1\epsilon_2} + \Delta_2 + \Delta_4 - \Delta_1 - \Delta_3 \quad (135)$$

(this rule can look more natural, if one mentions that in the free field model there is an identity $2\vec{\alpha}_2\vec{\alpha}_4 + \Delta_4 + \Delta_2 - \Delta_1 - \Delta_3 = 2\vec{\alpha}_1\vec{\alpha}_3$). This same permutation with the corresponding change of $\nu \rightarrow \frac{2S(\alpha_2)S(\alpha_4)}{\epsilon_1\epsilon_2} - \Delta_2 - \Delta_4 + \Delta_1 + \Delta_3$ was discussed in the Liouville case in [4].

With these values of parameters

$$\boxed{B^{(1)} = Z^{(1)} + \nu \quad \text{and} \quad B^{(2)} = Z^{(2)} + \nu Z^{(1)} + \frac{\nu(\nu+1)}{2},} \quad (136)$$

where Z and B are explicitly defined in (100) and (10) with substituted (64), (65) and (111). Of many formulas in this paper these five are the only ones used in the actual calculation at levels one and two (certainly, to check (136) one also needs the identifications of parameters in (130)-(134)). All the rest is included in order to present the logic and details of from-the-first-principles calculations very explicitly, so that they can be straightforwardly validated and generalized.

Note that if formulas are rewritten in terms of the fundamental weights vectors λ_i from (46), they admit immediate generalization to arbitrary N . The $sl(N)$ conformal Toda model has $W^{(N)}$ symmetry, and the eigenvalues of its generators in α -parametrization are given by

$$\Delta = \frac{\vec{\epsilon}^2 - \vec{\alpha}^2}{\epsilon_1\epsilon_2}, \quad \dots, \quad w_{\vec{\alpha}}^{(N)} \sim \frac{1}{\sqrt{(-\epsilon_1\epsilon_2)^N}} \prod_{i=1}^N (\vec{\alpha}\vec{\lambda}_i) \quad (137)$$

where $\vec{\epsilon} = \epsilon\vec{\rho}$ is directed along $\vec{\rho}$, the half-sum of all positive roots (or the sum of the fundamental weights), the central charge is $c = (N-1)\left(1 + N(N+1)\frac{\epsilon^2}{\epsilon_1\epsilon_2}\right)$ and $\{\vec{\lambda}_i\}$ is a Weyl-symmetric set of N minimal weights, see Fig.1. Then the AGT relation between the $sl(N)$ Toda and $U(N)$ Nekrasov's functions should look like

$$\boxed{a_i = (\sqrt{2}\vec{\alpha})\vec{\lambda}_i, \quad \mu_i^+ = -(\sqrt{2}\vec{\alpha}_2)\vec{\lambda}_i + S(\vec{\alpha}_1), \quad \mu_i^- = (\sqrt{2}\vec{\alpha}_4)\vec{\lambda}_i + S(\vec{\alpha}_3), \quad \nu = \frac{NS(\vec{\alpha}_1)S(\vec{\alpha}_3)}{\epsilon_1\epsilon_2}} \quad (138)$$

The $\sqrt{2}$ factors appear because with our normalization conditions the free field primaries are $V_{\vec{\alpha}} = e^{\sqrt{2}\vec{\alpha}\vec{\phi}}$. The notation S refers simultaneously to "shift" and "special".

Less technical is the general proof of the AGT relation. A first step is made in [6] and s.6 of the present paper. It concerns the case, when one of the two *special* Verma modules in the 4-point conformal block is *further* specialized by imposing two extra null-vector conditions which leads to additional selection rules on the intermediate state (enforced by the vanishing of the structure constants $C_{\alpha_1, \vec{\alpha}_2, \vec{\alpha}}$, which are not included into the definition of conformal block in the present text). In this particular case, the conformal block was explicitly found in [17] and it represents the generic hypergeometric function ${}_N F_{N-1}$, which has a straightforward character expansion. As rightly anticipated in [2], comparing it with Nekrasov formulas, which in this case get contribution only from a single set of "chiral" N -plets of Young diagrams, one obtains a complete *proof* of the AGT relation in this particular case. See [6] and s.6 above.

The AGT relations can provide a new breath to abandoned areas of conformal theory. The beautiful theory of W -algebras, for example, may not attract enough attention, because there was nothing to compare it with. Now *the equation gets the other side*: whatever one obtains by hard calculations in W theory can be compared with the differently looking formulas on the Nekrasov side. We demonstrated in this paper that this, indeed, opens a possibility of validating and improving W -algebra calculations.

As emphasized in [4], the AGT relation implies that there are two different expansions of the same quantity in characters (in sums over the Young diagrams): the natural expansion of conformal blocks and the Nekrasov sum over integer partitions. Already for $SU(2)$ they look absolutely different, like expansions associated with free boson and free fermion formalisms. The difference becomes even more pronounced for $N > 2$: it is enough to say that the conformal block expansion is in terms of generalized Young diagrams $\mathcal{Y}$, labeling elements of the W_N -algebra Verma modules, while the Nekrasov functions are labeled by N -ples of the ordinary Young diagrams, and there is no any *a priori* obvious relation between the two. It can deserve mentioning that the AGT relation does not link contributions of individual diagrams on the l.h.s. and the r.h.s., only the entire sums

over all diagrams of a given size (level). Thus, at the same stage it can be useful to consider relations between the conformal blocks of a given level and the original integrals from [8] which were afterwards decomposed into sums of the Nekrasov functions. **The AGT relation implies that these integrals have two different complementary combinatorial expansions** (we use the word combinatorial, like in [4], to emphasize that the character structure is not yet explicitly revealed in these expansions).

A real mystery at the moment is what substitutes the AGT relations in the case of generic Verma modules of the W_N algebras. The question can be asked in two opposite directions, that is, about more *general* and more *special* cases: what happens to generic *non-degenerate* modules for $N > 2$ and what happens to *stronger* degenerate modules (say, associated with the Ashkin-Teller model [31] and alike), which have more sophisticated conformal blocks, involving elliptic theta-constants already for $N = 2$. Answers to these questions can help to better understand the group theory meaning and possible generalizations of the Nekrasov functions.

The last but not least: within the general framework, **the AGT relation between conformal theories and Nekrasov functions looks very much like a "quantization" of the well-known link [32] between Seiberg-Witten theory [33] and integrable systems**. It is very important to understand the AGT relation from this perspective and to compare it with the other quantizations of the same link, like those of [34]-[36].

Appendix: Normalization conventions

We use somewhat non-standard normalizations, adjusted to maximally simplify the W -algebra formulas. The price for this is certain deviations from choices made in some other papers. We illustrate our conventions with the example of free field theory.

The freedom in the choice of normalization prescriptions in CFT can be described as follows. First, one can choose the coefficient in the action, i.e. normalization of the free field, or, equivalently, the coefficient in the propagator, we call it κ . If the stress tensor is defined as a generator of (local) translations, i.e. has the standard operator product expansion with the term $T(z_1)V(z_2) = \dots + z_{12}^{-1}\partial V(z_2) + \dots$ with the *unit* coefficient in front of the V -derivative term, then, κ^{-1} appears as the coefficient in T , thus, κ can be alternatively considered as a freedom in normalization of $T = W^{(2)}$. Similarly, to each operator $W^{(k)}$ (there are $N - 1$ of them in the theory with $N - 1$ free fields) can be ascribed an arbitrary new coefficient κ_k , i.e. there is no way to tie the normalization of $W^{(k)}$'s with $k > 2$ to that of the propagator, and there is no *canonical* way to fix normalizations of some structure constants of the W -algebra. When the central charge c deviates from $N - 1$, there is a new arbitrariness in the choice of the normalization parameter in the deformation term $\partial^2\phi$ in T : in fact, it can be considered as a freedom to normalize $W^{(1)}$. Second, in addition to these N free normalizations, different papers define differently the parameter λ in the conformal dimensions: $\Delta_{\vec{\alpha}} = \Delta(e^{\vec{\alpha}\phi/\lambda})$. So far, fortunately the same λ always appeared in the definition of other eigenvalues: $w_{\vec{\alpha}} = w(e^{\vec{\alpha}\phi/\lambda})$ and so on.

More explicitly, if the operator product expansion of the free fields is defined as

$$\partial\phi_i(z)\partial\phi_j(0) = \kappa_2 \frac{\delta_{ij}}{z^2} + \dots \quad (139)$$

then the stress tensor

$$T(z) = \frac{1}{2\kappa_2} \sum_{i=1}^{N-1} (\partial\phi_i)^2 + \kappa_1 \sum_{i=1}^{N-1} Q_i \partial^2\phi_i = \frac{1}{2\kappa_2} (\partial\vec{\phi})^2 + \kappa_1 \vec{Q} \partial^2\vec{\phi} \quad (140)$$

satisfies

$$T(z)T(0) = \frac{c/2}{z^4} + \frac{2T(0)}{z^2} + \frac{\partial T(0)}{z} + \dots \quad (141)$$

with

$$c = N - 1 - 12\kappa_2\kappa_1^2\vec{Q}^2, \quad (142)$$

while the primary field

$$V_{\vec{\alpha}} = : \exp\left(\frac{\vec{\alpha}\vec{\phi}}{\lambda}\right) : \quad (143)$$

satisfies

$$T(z)V_{\vec{\alpha}}(0) = \frac{\Delta_{\vec{\alpha}}}{z^2} V_{\vec{\alpha}}(0) + \frac{1}{z} L_{-1} V_{\vec{\alpha}}(0) + \dots \quad (144)$$

with the dimension

$$\Delta_{\vec{\alpha}} = \frac{\kappa_2}{\lambda^2} \cdot \frac{\vec{\alpha}(\vec{\alpha} - 2\lambda\kappa_1\vec{Q})}{2} \quad (145)$$

The free field theory selection rule for the non-vanishing correlator of exponentials is

$$\sum_i \vec{\alpha}_i = 2\lambda\kappa_1\vec{Q} = 2\lambda\varepsilon\vec{\rho} \quad (146)$$

It is inspired by zero-mode integration within the functional integral approach, or by conditions like

$$\langle V_{\vec{\alpha}_1}(z_1)V_{\vec{\alpha}_2}(z_2) \rangle \sim (z_1 - z_2)^{\kappa_2/\lambda^2} = \frac{1}{(z_1 - z_2)^{\Delta_{\vec{\alpha}_1} + \Delta_{\vec{\alpha}_2}}} \implies \Delta_{\vec{\alpha}_1} = \Delta_{\vec{\alpha}_2} = -\frac{\kappa_2}{2\lambda^2} \vec{\alpha}_1 \vec{\alpha}_2 \quad (147)$$

within CFT itself.

Usually in CFT with $N \neq 2$ [12] one chooses $\vec{Q}$ to be directed along the vector $\vec{\rho}$, the half-sum of all positive roots of $sl(N)$ or, which is the same, the sum of all fundamental weights, $|\vec{\rho}|^2 = \frac{N(N^2-1)}{12}$ (for non-simply-laced groups $\vec{Q} \sim \sqrt{-\epsilon_1/\epsilon_2}\vec{\rho} + \sqrt{-\epsilon_2/\epsilon_1}\vec{\rho}^\vee$). The proportionality coefficient, however, is not canonically fixed and remains an arbitrary normalization parameter. The apparent problem with this convention, uniform and convenient for all groups, is that for $SU(2)$ $\rho = 1/\sqrt{2}$, therefore, one rarely uses it in considerations of the single field case, since it makes k_1 to contain $\sqrt{2}$. One can continue this "natural" $N = 2$ convention to all N , using coordinates where $\vec{\rho}\vec{\phi}/|\rho| = \phi_{N-1}$, and this choice also is present in the literature. Anyhow,

$$\boxed{\text{if } \kappa_1\vec{Q} = \varepsilon\vec{\rho}, \quad \text{then } c = (N-1)\left(1 - N(N+1)\kappa_2\varepsilon^2\right) \quad \text{and} \quad \Delta_{\vec{\alpha}} = \frac{\kappa_2}{2\lambda^2} \vec{\alpha}(\vec{\alpha} - 2\lambda\kappa_1\vec{Q})} \quad (148)$$

At the same time, the central charge of the $sl(N)$ Toda model, expressed by the AGT relation through the Nekrasov parameters $\epsilon = \epsilon_1 + \epsilon_2$, is equal to

$$\boxed{c_N = (N-1) \left(1 + N(N+1) \frac{\epsilon^2}{\epsilon_1\epsilon_2}\right) = \begin{cases} 1 + \frac{6\epsilon^2}{\epsilon_1\epsilon_2} & \text{for } N = 2, \\ 2 + \frac{24\epsilon^2}{\epsilon_1\epsilon_2} & \text{for } N = 3, \\ \dots & \end{cases}} \quad (149)$$

i.e. in this notation

$$\varepsilon = \frac{1}{\sqrt{\kappa_2}} \cdot \frac{\epsilon}{\sqrt{-\epsilon_1\epsilon_2}} \quad (150)$$

Since the common rescaling of all parameters in the Nekrasov functions with $N_f = 2N$ does not affect the answer, one can easily choose $\epsilon_2 = -1/\epsilon_1$, as it was done, for example, in [1] and [6]. The following table contains the comparison of notations in different papers:

paper	N	central charge	dimension	selection rule : $\sum_i \vec{\alpha}_i =$	exponential primary	$1/\lambda$	$\varepsilon/Q = \kappa_1/ \rho $	$\kappa_1 \vec{Q}$	κ_2
[2]	2	$1 + 6Q^2$	$\alpha(Q - \alpha)$	Q	$: e^{2\alpha\phi} :$	2	$\sqrt{2}$	Q	$-1/2$
[2]	N	$(N-1)\left(1 + \frac{1}{N(N+1)\varepsilon^2}\right)$	$\frac{\vec{\alpha}(2\vec{Q}-\vec{\alpha})}{2}$	$2\vec{Q}$	$: e^{\vec{\alpha}\vec{\phi}} :$	1	$\frac{1}{ \rho } = \sqrt{\frac{12}{N(N^2-1)}}$	$\vec{Q}$	-1
[4]	2	$1 + \frac{6Q^2}{\varepsilon_1\varepsilon_2}$	$\frac{\alpha(Q-\alpha)}{\varepsilon_1\varepsilon_2}$	Q	$: e^{\alpha\phi} :$	1	$\frac{1}{2\sqrt{2}}$	$\frac{1}{2}Q$	$\frac{2}{\varepsilon_1\varepsilon_2}$
[5]	2	$1 - 12Q^2$	$\frac{\alpha(\alpha-2Q)}{2}$	$2Q$	$: e^{\alpha\phi} :$	1	$ \rho_{SU(2)} ^{-1} = \sqrt{2}$	Q	1
[5]	3	$2 - 12Q^2$	$\frac{\alpha^2 + \beta(\beta-2Q)}{2}$	$(0, 2Q)$	$: e^{\alpha\phi_1 + \beta\phi_2} :$	1	$ \rho_{SU(3)} ^{-1} = \frac{1}{\sqrt{2}}$	$(0, Q)$	1
present paper	3	$2 - 24Q^2$	$\alpha^2 + \tilde{\beta}^2 - Q^2$ $\tilde{\beta} = \beta - Q$	$(0, 2Q)$	$: e^{\sqrt{2}(\alpha\phi_1 + \beta\phi_2)} :$	$\sqrt{2}$	1	$(0, \sqrt{2}Q)$	1
[6] and present paper	N	$(N-1)\left(1 + \frac{N(N+1)\varepsilon^2}{\varepsilon_1\varepsilon_2}\right)$	$\frac{\vec{\alpha}(\vec{\alpha}-2\vec{Q})}{-\varepsilon_1\varepsilon_2}$ $= \frac{\vec{\alpha}^2 - Q^2}{-\varepsilon_1\varepsilon_2}$	$2\vec{Q}$	$: e^{\sqrt{2}\vec{\alpha}\vec{\phi}} :$	$\sqrt{2}$	$\frac{\sqrt{2}}{ \rho }$	$\sqrt{2}\vec{Q}$	$\frac{1}{-\varepsilon_1\varepsilon_2}$

In sample free field considerations with $N = 3$ in this paper we imply that $V_{\vec{\alpha}} = e^{\sqrt{2}(\alpha\phi_1 + \beta\phi_2)}$. Thus, it is natural to normalize the $W^{(3)}$ operator to be

$$W^{(3)} = \frac{1}{2\sqrt{2}} \left((\partial\phi_1)^3 - 3\partial\phi_1(\partial\phi_2)^2 \right) \quad (151)$$

so that

$$W^{(3)}(z)V_{\vec{\alpha}}(0) = \frac{w_{\vec{\alpha}}}{z^3}V_{\vec{\alpha}}(0) + \frac{1}{z^2}W_{-1}V_{\vec{\alpha}}(0) + \frac{1}{z}W_{-2}V_{\vec{\alpha}}(0) \dots \quad (152)$$

with⁵

$$w_{\vec{\alpha}} = \alpha(\alpha^2 - 3\beta^2) \quad (153)$$

and

$$W_{-1}V_{\vec{\alpha}} = : \frac{3}{\sqrt{2}} \left((\alpha^2 - \beta^2)\partial\phi_1 - 2\alpha\beta\partial\phi_2 \right) V_{\vec{\alpha}} : \quad (154)$$

Also, of course,

$$L_{-1}V_{\vec{\alpha}} = : \sqrt{2}(\alpha\partial\phi_1 + \beta\partial\phi_2)V_{\vec{\alpha}} : = \partial V_{\vec{\alpha}} \quad (155)$$

The *special* primaries of the W_3 algebra generate the Verma modules with a null-vector at the first level, with $W_{-1}V_{\vec{\alpha}} = \zeta L_{-1}V_{\vec{\alpha}}$ for some λ . From (155) and (154) it follows that

$$\begin{aligned} 3(\alpha^2 - \beta^2) &= 2\zeta\alpha, \\ -6\alpha\beta &= 2\zeta\beta \end{aligned} \quad (156)$$

⁵In fact, it is $w_{\vec{\alpha}} = (-\varepsilon_1\varepsilon_2)^{-3/2} \cdot \alpha(\alpha^2 - 3\beta^2)$, but we systematically suppress powers of $(-\varepsilon_1\varepsilon_2)$ in most formulas to make them more readable. One should remember they are actually there.

i.e. $\zeta = -3\alpha$ and $\beta^2 = 3\alpha^2$, or $\zeta = 3/2\alpha$ and $\beta = 0$. Note that, under these relations, $\zeta = 3w_{\vec{\alpha}}/2\Delta_{\vec{\alpha}}$, so that the null-vector condition is exactly (4) and the Kac determinant $v^2 = \Delta^3 - w^2 = \left(\beta(\beta^2 - 3\alpha^2)\right)^2$ vanishes.

The main peculiarity of our normalization conventions is that the dimension $\Delta = \vec{\alpha}^2$ does not contain a minus sign, in variance from [1, 2, 4]: in dealing with W -algebras this eliminates unnecessary imaginary units. Factors of 2 are eliminated from Δ and w to make the central relation (9) as simple as possible. The price for this is the roots of 2 in the primary exponentials and W operators. Roots of 3, however, are unavoidable, they come from the roots and weights of the underlying $sl(3)$ algebra.

Acknowledgements

We are indebted for hospitality and support to Prof.T.Tomas and the Institute of Theoretical and Computational Physics of University of Crete, where part of this work was done.

The work was partly supported by Russian Federal Nuclear Energy Agency and by RFBR grants 07-02-00878 (A.Mir.), and 07-02-00645 (A.Mor.). The work was also partly supported by grant FP7-REGPOT-1 (Crete HEP Cosmo 228644), by joint grants 09-02-90493-Ukr, 09-02-93105-CNRS, 09-01-92440-CE, 09-02-91005-ANF, INTERREG IIIA (Greece-Cyprus) and by Russian President's Grant of Support for the Scientific Schools NSH-3035.2008.2.

References

- [1] L.Alday, D.Gaiotto and Y.Tachikawa, arXiv:0906.3219
- [2] N.Wyllard, arXiv:0907.2189
- [3] D.Gaiotto, arXiv:0908.0307
- [4] A.Marshakov, A.Mironov and A.Morozov, arXiv:0907.3946
- [5] Andrey Mironov, Sergey Mironov, Alexei Morozov and Andrey Morozov, arXiv:0908.2064
- [6] A.Mironov and A.Morozov, arXiv:0908.2190
- [7] A.Belavin, A.Polyakov, A.Zamolodchikov, Nucl.Phys., **B241** (1984) 333-380
- [8] N.Nekrasov, Adv.Theor.Math.Phys. **7** (2004) 831-864
R.Flume and R.Pogossian, Int.J.Mod.Phys. **A18** (2003) 2541
N.Nekrasov and A.Okounkov, hep-th/0306238
- [9] A.Zamolodchikov, Comm.Math.Phys. **96** (1984) 419; Theor.Math.Phys. **73** (1987) 1088
- [10] A.Zamolodchikov and A.Zamolodchikov, *Conformal field theory and critical phenomena in 2d systems*, 2009, 168 p. (in Russian)
P.Di Francesco, P.Mathieu and D.Sénéchal, *Conformal Field Theory*, Springer, 1997
- [11] H.Dorn, H.-J.Otto, Phys.Lett., **B291** (1992) 39-43, hep-th/9206053; Nucl.Phys., **B429** (1994) 375-388, hep-th/9403141
A.Zamolodchikov and A.Zamolodchikov, Nucl.Phys. **B477** (1996) 577-605, hep-th/9506136
- [12] V.Dotsenko and V.Fateev, Nucl.Phys., **B240** (1984) 312-348
B.Feigin and D.Fuks, Funct.Anal.Appl. **16** (1982) 114-126 (Funkts.Anal.Pril. **16** (1982) 47-63)
A.Gerasimov, A.Marshakov, A.Morozov, M.Olshanetsky and S.Shatashvili, Int.J.Mod.Phys., **A5** (1990) 2495-2589
A.Gerasimov, A.Marshakov and A.Morozov, Nucl. Phys. **B328** (1990) 664-676
- [13] A.Zamolodchikov, Theor.Math.Phys. **65** (1985) 1205-1213 (Teor.Mat.Fiz. **65** (1985) 347-359)
V.Fateev and A.Zamolodchikov, Nucl.Phys. **B280** (1987) 644-660
- [14] V.Fateev and S.Lukyanov, Int.J.Mod.Phys. **A3** (1988) 507
- [15] A.Marshakov and A.Morozov, Nucl.Phys. **B339** (1990) 79-94
- [16] P. Bowcock and G.M.T. Watts, Theor.Math.Phys. **98** (1994) 350-356 (Teor.Mat.Fiz. **98** (1994) 500-508), arXiv:hep-th/9309146

- [17] V.Fateev and I.Litvinov, JHEP **0711** (2007) 002, arXiv:0709.3806
- [18] H.Sonoda, Nucl.Phys. **B311** (1988/89) 401-416
- [19] G.Moore and N.Seiberg, Comm.Math.Phys. **123** (1989) 177-254
- [20] E.Bergshoeff, C.Pope, L.Romans, E.Sezgin, X.Shen and K.Stelle, Phys.Lett. **B243** (1990) 350
 G.Sotkov and M.Stanishkov, Nucl.Phys. **B356** (1991) 439
 G.Sotkov, M.Stanishkov and C.J.Zhu, Nucl.Phys. **B356** (1991) 245
 A.Gerasimov, A.Levin and A.Marshakov, Nucl.Phys. **B360** (1991) 537
 A.Bilal, V.Fock and I.Kogan, Nucl.Phys. **B359** (1991) 635
 P.Mansfield and B.Spence, Nucl.Phys. **B362** (1991) 294
 J.-L.Gervais and Y.Matsuo, Preprint LPTENS-91/35, NBI-HE-91-50
 C. Itzykson, “*W* - geometry,” Cargese lectures, appeared in *Random surfaces and Quantum gravity*, ed. O. Alvarez *et al.*, Plenum press, New York, 1991
 A.Marshakov, A.Mironov, A.Morozov and M.Olshanetsky, Nucl.Phys. **B404** (1993) 427-456, hep-th/9203044
 J. Gomis, J. Herrero, K. Kamimura and J. Roca, Phys.Lett. **B339** (1994) 59-64, hep-th/9409024
 S.Govindarajan, Nucl.Phys. **B457** (1995) 357-374, hep-th/9412078
 M.Olshanetsky, nlin/0011010
- [21] D.Gaiotto, arXiv:0904.2715
 See also:
 Y.Tachikawa, JHEP **0907** (2009) 067, arXiv:0905.4074
 D.Nanopoulos and D.Xie, arXiv:0907.1651
- [22] E.Witten, Nucl.Phys., **B500** (1997) 3-42, hep-th/9703166
- [23] A.Marshakov, M.Martellini and A.Morozov, Phys.Lett. **B418** (1998) 294-302, hep-th/9706050
 A.Gorsky, S.Gukov and A.Mironov, Nucl.Phys., **B517** (1998) 409-461, hep-th/9707120; Nucl.Phys., **B518** (1998) 689, hep-th 9710239
- [24] A. Mironov, A. Morozov and L. Vinet, Teor.Mat.Fiz. **100** (1994) 119-131 (Theor.Math.Phys. **100** (1995) 890-899), hep-th/9312213;
 A.Gerasimov, S.Khoroshkin, D.Lebedev, A.Mironov and A.Morozov, Int.J.Mod.Phys. **A10** (1995) 2589-2614, hep-th/9405011;
 S.Kharchev, A.Mironov and A.Morozov, q-alg/9501013;
 A.Mironov, hep-th/9409190; Theor.Math.Phys. **114** (1998) 127, q-alg/9711006
- [25] A.Alexandrov, A.Mironov and A.Morozov, Int.J.Mod.Phys. **A19** (2004) 4127, Theor.Math.Phys. **142** (2005) 349, hep-th/0310113; Physica **D235** (2007) 126-167, hep-th/0608228; hep-th/0605171;
- [26] V.Schechtman and A.Varchenko, *Integral representations of N-point conformal correlators in the WZW model*, Preprint MPI/89-51, Max Planck Institut, Bonn(1989), 1-21
- [27] A.Morozov and L.Vinet, Mod.Phys.Lett. **A8** (1993) 2891-2902, hep-th/9111023
- [28] D.E.Littlewood, *The theory of group characters and matrix representations of groups*, Oxford, 1958
 M.Hamermesh, *Group theory and its application to physical problems*, 1989
 I.G.Macdonald, *Symmetric functions and Hall polynomials*, Oxford Science Publications, 1995
 W.Fulton, *Young tableaux: with applications to representation theory and geometry*, London Mathematical Society, 1997
- [29] S.Kharchev, A.Marshakov, A.Mironov and A.Morozov, Int.J.Mod.Phys., **A10** (1995) 2015-2052, hep-th/9312210
 A.Mironov, A.Morozov and G.Semenoff, Int.J.Mod.Phys., **A10** (1995) 2015, hep-th/9404005
 I.Kostov, M.Staudacher and T.Wynter, Commun.Math.Phys. **191** (1998) 283-298, hep-th/9703189
 A.Morozov and Sh.Shakirov, arXiv: 0902.2627
 A.Mironov, A.Morozov and S.Natanzon, arXiv:0904.4227
 A.Morozov, arXiv:0906.3518
- [30] R.Dijkgraaf, In: *The moduli spaces of curves*, Progress in Math., 129 (1995), 149-163, Birkhäuser
 I.Goulden and D.Jackson, Proc.Amer.Math.Soc. **125** (1997) 51-60, math/9903094
 A.Givental, math/0108100

T.Ekedahl, S.Lando, M.Shapiro, A.Vainshtein, Invent.Math.146(2001),297-327
 S.Lando, Russ.Math.Surv., **57** (2002) 463-533
 M.Kazarian and S.Lando, math.AG/0410388; math/0601760
 M.Kazarian, arXiv:0809.3263
 V.Bouchard and M.Marino, arXiv:0709.1458
 A.Mironov and A.Morozov, JHEP **0902** (2009) 024, arXiv:0807.2843

- [31] The first review in [10]
 Al.Zamolodchikov, Nucl.Phys. **B285** [**FS19**] (1987) 481
 M.Bershadsky and A.Radul, Int.J.Mod.Phys. **A2** (1987)165
 L.J.Dixon, D.Friedan, E.J.Martinec and S.H.Shenker, Nucl.Phys. **B282** (1987) 13
 H.La, BUHEP-88-37
 D.Lebedev and A.Morozov, Nucl.Phys. **B302** (1988) 163
- [32] A.Gorsky, I.Krichever, A.Marshakov, A.Mironov, A.Morozov, Phys.Lett., **B355** (1995) 466-477
 R.Donagi and E.Witten, Nucl.Phys., **B460** (1996) 299-334
 E.Martinec, Phys.Lett., **B367** (1996) 91-96, hep-th/9510204
 A.Gorsky, A.Marshakov, Phys.Lett., **B374** (1996) 218-224, hep-th/9510224
 H.Itoyama and A.Morozov, Nucl.Phys., **B477** (1996) 855-877, hep-th/9511126; Nucl.Phys., **B491** (1997) 529-573, hep-th/9512161
 A.Gorsky, A.Marshakov, A.Mironov and A.Morozov, Phys.Lett., **B380** (1996) 75-80, arXiv:hep-th/9603140
 A.Gorsky and A.Mironov, hep-th/0011197
- [33] N.Seiberg and E.Witten, Nucl.Phys., **B426** (1994) 19-52; Nucl.Phys., **B431** (1994) 484-550
- [34] A.Gorsky, A.Marshakov, A.Mironov and A.Morozov, Nucl.Phys., **B527** (1998) 690-716, hep-th/9802004
 J.Edelstein, M.Marino and J.Mas, Nucl.Phys., **B541** (1999) 671-697, hep-th/9805172
 J.Edelstein, M.Gomez-Reino and M.Marino, Adv.Theor.Math.Phys., **4** (2000) 503-543, hep-th/0006113
- [35] R.Dijkgraaf and C.Vafa, Nucl.Phys. **B644** (2002) 3, hep-th/0206255; Nucl.Phys. **B644** (2002) 21, hep-th/0207106; hep-th/0208048
 L.Chekhov and A.Mironov, Phys.Lett. **B552** (2003) 293, hep-th/0209085
 R.Dijkgraaf, S.Gukov, V.Kazakov and C.Vafa, Phys.Rev. **D68** (2003) 045007, hep-th/0210238
 V.Kazakov and A.Marshakov, J.Phys. **A36** (2003) 3107-3136, hep-th/0211236
 H.Itoyama and A.Morozov, Nucl.Phys. **B657** (2003) 53, hep-th/0211245; Phys.Lett. **B555** (2003) 287, hep-th/0211259; Prog.Theor.Phys. **109** (2003) 433, hep-th/0212032; Int.J.Mod.Phys. **A18** (2003) 5889, hep-th/0301136
 S.Naculich, H.Schnitzer and N. Wyllard, JHEP **0301** (2003) 015, hep-th/0211254;
 B.Feng, Nucl.Phys. **B661** (2003) 113-138, hep-th/0212010;
 I.Bena, S.de Haro and R.Roiban, Nucl.Phys. **B664** (2003) 45-58, hep-th/0212083;
 Ch.Ann, Phys.Lett. **B560** (2003) 116-127, hep-th/0301011;
 L.Chekhov, A.Marshakov, A.Mironov and D.Vasiliev, hep-th/0301071; Proc. Steklov Inst.Math. **251** (2005) 254, hep-th/0506075
 A.Mironov, Theor.Math.Phys. **146** (2006) 63-72, hep-th/0506158;
 A. Dymarsky and V. Pestun, Phys.Rev. **D67** (2003) 125001, hep-th/0301135
- [36] N.Nekrasov and S.Shatashvili, arXiv:0901.4748